

Assignment #1

Due: Friday, February 11, 2005

Consider the following questions for Na. Try to plug in numbers after obtaining an analytical expression.

- Na oven temperature $T = 600K$
- natural line width $\gamma = 2\pi \times 10$ MHz
- Zeeman splitting 1.4 MHz/gauss
- wavelength $\lambda = 2\pi/k = 589$ nm

1. A Zeeman Slower

If you want to slow an atomic beam efficiently, you have to compensate for the changing Doppler shift ($\vec{k} \cdot \vec{v}$) during the deceleration. This can be done by sweeping the frequency of the slowing beam, i.e., chirping (as discussed in class). A method of producing a continuous beam of cold atoms is Zeeman slowing.

A well collimated beam of atoms is originating from an oven with a temperature T . The beam propagates along a distance L with a longitudinal magnetic field $B(x)$ ($0 < x < L$). A laser beam of intensity I is counter propagating. Its frequency is detuned by δ ($\delta \equiv \omega - \omega_0$) from the transition frequency at $B = 0$.

- a) Calculate the maximum deceleration a_{max} you can achieve. Assume you could choose arbitrarily large laser intensities.

Assume you want to slow down atoms with speeds lower than the peak velocity v_{max} of the thermal distribution to a stand still using the constant deceleration $f a_{max}$. ($0 < f < 1$) The Zeeman effect shifts the resonance $\omega_0 \rightarrow \omega_0 + g\mu_B B(x)$ (where $g\mu_B = (2\pi)1.4$ MHz/gauss, in this case).

- b) Calculate the spatial dependence of the magnetic field $B(x)$ and the length L of the slower.
- c) Assume three different models for spontaneous emission: (i) Photons are emitted along the $+/ -\hat{x}$ direction, (ii) isotropic emission, or (iii) emission in a dipole pattern. What is the momentum diffusion constant \mathcal{D} for the (longitudinal) x-component of the momentum in the three cases, at a given photon scattering rate Γ_s ?

2. Slowing an atom with off-resonant light

Assume you want to slow an atom of velocity v_{max} with a counter propagating laser beam that is on resonance with the atom at rest. (Use the same v_{max} as in Problem 1, and $I = 5I_{sat}$.)

- a) How long would it take ?
- b) How far would the atom travel ?

hint: You can easily integrate the equation of motion

3. Classical molasses in a vapor cell

Assume you have an experimental setup with 3 dimensional molasses in a spherical volume of radius $r = 2mm$. The molasses is surrounded by a gas with pressure $p = 10^{-8}$ torr and a temperature $T = 350K$.

- a) Estimate the velocity capture range v_{cap} of this classical molasses. (v_{cap} is the velocity of the fastest atom that can be captured.) There are six laser beams, each with intensity $I = I_{sat}/6$. (hint: use an approximate expression for the force for $v \gg \gamma/k$)
- b) Choose the detuning and the laser intensity in order to maximize the diffusion time in the molasses. Calculate the maximum diffusion time.