

MIT Open Access Articles

ROSSITER-MCLAUGHLIN OBSERVATIONS OF 55 Cnc e

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: López-Morales, Mercedes, Amaury H. M. J. Triaud, Florian Rodler, Xavier Dumusque, Lars A. Buchhave, Avet Harutyunyan, Sergio Hoyer, et al. "ROSSITER-MCLAUGHLIN OBSERVATIONS OF 55 Cnc e." *The Astrophysical Journal* 792, no. 2 (August 25, 2014): L31. © 2014 American Astronomical Society.

As Published: <http://dx.doi.org/10.1088/2041-8205/792/2/l31>

Publisher: Institute of Physics/American Astronomical Society

Persistent URL: <http://hdl.handle.net/1721.1/93251>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



ROSSITER–MCLAUGHLIN OBSERVATIONS OF 55 Cnc e

MERCEDES LÓPEZ-MORALES¹, AMAURY H. M. J. TRIAUD^{2,15}, FLORIAN RODLER^{1,3,16}, XAVIER DUMUSQUE¹,
LARS A. BUCHHAVE^{1,4}, AVET HARUTYUNYAN⁵, SERGIO HOYER^{6,7,17}, ROI ALONSO^{6,7}, MICHAËL GILLON⁸, NATHAN A. KAIB⁹,
DAVID. W. LATHAM¹, CHRISTOPHE LOVIS¹⁰, FRANCESCO PEPE¹⁰, DIDIER QUELOZ^{10,11}, SEAN N. RAYMOND^{12,13},
DAMIEN SÉGRANSAN¹⁰, INGO P. WALDMANN¹⁴, AND STÉPHANE UDRY¹⁰

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 01238, USA; mlopez-morales@cfa.harvard.edu

² Department of Physics, and Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

³ Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

⁴ Centre for Star and Planet Formation, Natural History Museum of Denmark, University of Copenhagen, DK-1350 Copenhagen, Denmark

⁵ Fundación Galileo Galilei-INAf, Rambla José Ana Fernández Pérez, E-738712 Breña Baja, Tenerife, Spain

⁶ Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain

⁷ Departamento de Astrofísica, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Spain

⁸ Institut d’Astrophysique et Géophysique, Université de Liège, Allée du 6 Août 17, B-4000 Liège, Belgium

⁹ Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA) and Department of Physics and Astronomy, Northwestern University, 2131 Tech Drive, Evanston, IL 60208, USA

¹⁰ Observatoire Astronomique de l’Université de Genève, Chemin des Maillettes 51, Sauverny, CH-1290, Switzerland

¹¹ Cavendish Laboratory, J J Thomson Avenue, Cambridge CB3 0HE, UK

¹² Université de Bordeaux, LAB, UMR 5804, BP F-33270 Floirac, France

¹³ CNRS, LAB, UMR 5804, F-33270 Floirac, France

¹⁴ Department of Physics and Astronomy, University College London, Gower Street, WC1E6BT, UK

Received 2014 June 27; accepted 2014 August 8; published 2014 August 25

ABSTRACT

We present Rossiter–McLaughlin observations of the transiting super-Earth 55 Cnc e collected during six transit events between 2012 January and 2013 November with HARPS and HARPS-N. We detect no radial velocity signal above 35 cm s^{-1} (3σ) and confine the stellar $v \sin i_*$ to $0.2 \pm 0.5 \text{ km s}^{-1}$. The star appears to be a very slow rotator, producing a very low amplitude Rossiter–McLaughlin effect. Given such a low amplitude, the Rossiter–McLaughlin effect of 55 Cnc e is undetected in our data, and any spin–orbit angle of the system remains possible. We also performed Doppler tomography and reach a similar conclusion. Our results offer a glimpse of the capacity of future instrumentation to study low amplitude Rossiter–McLaughlin effects produced by super-Earths.

Key words: planets and satellites: formation – planets and satellites: individual (55 Cancri e) – stars: individual (55 Cancri) – techniques: radial velocities – techniques: spectroscopic

Online-only material: color figures, machine-readable table

1. INTRODUCTION

55 Cnc is a $0.9 M_{\odot}$ star (von Braun et al. 2011) harboring five known planets with masses between $0.025 M_{\text{Jup}}$ and $3.8 M_{\text{Jup}}$, and orbital periods between ~ 0.74 days and ~ 4872 days (Nelson et al. 2014). The star also has a $\sim 0.27 M_{\odot}$ binary companion at a projected orbital separation of 1065 AU (Mugrauer et al. 2006).

The innermost planet in the system, 55 Cnc e ($M_p = 8.3 \pm 0.4 M_{\text{Earth}}$; $R_p = 1.94 \pm 0.08 R_{\text{Earth}}$), was found to transit by Winn et al. (2011) and Demory et al. (2011), after Dawson & Fabrycky (2010) provided a revised period of 0.74 days, shorter than the originally reported 2.8 days period by McArthur et al. (2004). The presence of transits makes 55 Cnc e an invaluable potential target for future studies of atmospheric properties of super-Earths and spin–orbit angle studies via the Rossiter–McLaughlin effect (Rossiter 1924; McLaughlin 1924; Queloz et al. 2000; Gaudi & Winn 2007). In the case of spin–orbit angle studies, a result for 55 Cnc e would add to the very few measurements reported so far for Neptune and super-Earth mass planets, e.g., the detection of oblique orbits for HAT-P-11b (Winn et al. 2010; Hirano et al. 2011), and for the

multiple super-Earth systems Kepler-50 and Kepler-65—these last two via asteroseismology (Chaplin et al. 2013). There is also the non-detection of the Rossiter–McLaughlin effect of GJ 436 by Albrecht et al. (2012), which indicates that the star is a very slow rotator with $v \sin i_* < 0.4 \text{ km s}^{-1}$.

Kaib et al. (2011) showed that the distant binary companion to 55 Cnc causes the planetary system to precess as a rigid body (see also Innanen et al. 1997; Batygin et al. 2011; Boué & Fabrycky 2014). The planets’ orbits nonetheless remain confined to a common plane such that a measurement of planet e’s spin–orbit angle should be representative of the entire system. Given the unknown orientation of the wide binary orbit, Kaib et al. (2011) calculated that the plane of the planets is most likely tilted with respect to the stellar equator. The plane could be tilted by virtually any angle, with a most probable projected spin–orbit angle of $\sim 50^\circ$. There is also a $\sim 30\%$ chance of a retrograde configuration.

Valenti & Fischer (2005) estimated a $v \sin i_*$ for 55 Cnc of $2.4 \pm 0.5 \text{ km s}^{-1}$, which, combined with the depth of the observed transit, was expected to yield a Rossiter–McLaughlin effect with a semi-amplitude of $70 \pm 15 \text{ cm s}^{-1}$. Such an amplitude should be detectable with stabilized, high-resolution spectrographs such as HARPS (Molaro et al. 2013).

Following those results, we set out to detect the Rossiter–McLaughlin effect of 55 Cnc e and test the spin–orbit misalignment prediction of the system by Kaib et al. (2011).

¹⁵ Swiss National Science Foundation Fellow.

¹⁶ Alexander-von-Humboldt Postdoctoral Fellow.

¹⁷ Severo Ochoa Fellow.

Table 1
Radial Velocities of all Spectra of 55 Cnc Observed with HARPS and HARPS-N

BJD	RV	σ_{RV}	FWHM	BIS _{span}	S/N	Airmass	Seeing	Exposure	RV _{cor}
−2400000	(km s ^{−1})	(km s ^{−1})	(km s ^{−1})	(km s ^{−1})	at 6240 Å	...	(arcsec)	(s)	(km s ^{−1})
55954.663365	27.42076	0.00090	6.22245	−0.01210	118.80	2.04	0.74	123.1	−0.00298
55954.665229	27.41805	0.00082	6.22627	−0.00865	133.50	2.03	0.81	123.1	0.00044
55954.667405	27.41877	0.00066	6.22736	−0.00984	181.50	2.01	0.78	180.0	−0.00121
55954.669905	27.41707	0.00066	6.22624	−0.01121	178.90	2.00	0.72	180.0	0.00044
55954.672474	27.41806	0.00065	6.22813	−0.01089	182.80	1.98	0.76	180.0	−0.00061
...

Notes. The columns are the following: barycentric Julian date, radial velocity (RV), RV errorbars, FWHM of the cross-correlation function, span in bisector slope as defined in Queloz et al. (2000), signal-to-noise ratio at 6240 Å, airmass, seeing measured by the La Silla seeing monitor, exposure time, and RVs with the five planet Doppler displacement removed.

(This table is available in its entirety in a machine-readable form in the online journal. A portion is shown here for guidance regarding its form and content.)

2. OBSERVATIONS

Shortly after the detection of 55 Cnc e’s transit was announced, we requested four spectroscopic time series on HARPS (Prog. ID 288.C-5010; PI: Triaud) as Director Discretionary Time. HARPS is installed on the 3.6 m telescope at the European Southern Observatory on La Silla, Chile (Mayor et al. 2003). The position of 55 Cnc in the sky—R.A.(J2000) = 08:52:35.81, decl.(J2000) = +28:10:50.95—is low as seen from La Silla. The target remains at a zenith distance of $z < 2$ for only ~ 2.5 hr per night, with a transit duration of about 1.5 hr having to fit within this tight window. This constraint on the airmass, essential to obtain precise radial velocities (RVs), is set by the instrumental atmospheric dispersion corrector. We used the ephemeris by Gillon et al. (2012), then at an advanced stage of preparation, to schedule our observations. In total, we gathered 179 spectra on the nights starting on 2012 January 27, 2012 February 13, 2012 February 27, and 2012 March 15 UT.

The target is better suited for observations from the north and, therefore, in 2013 we acquired additional RV time series with the newly installed HARPS-N instrument on the 3.57 m Telescopio Nazionale Galileo (TNG) at the Observatorio del Roque de los Muchachos in La Palma, Spain (Cosentino et al. 2012). HARPS-N is an updated version of HARPS and is able to reach the same overall RV precision, if not better (see, e.g., Pepe et al. 2013; Dumusque et al. 2014). The HARPS-N time was awarded via the Spanish TAC (Prog. ID 34-TNG4/13B; PI: Rodler) and observations were collected on the nights starting on 2013 November 14 and 2013 November 28 UT. We gathered a total of 113 spectra in those two nights. A third night awarded to this program was lost to weather. Some more details about the observations are provided in Table 1.

3. RADIAL VELOCITY EXTRACTION

The spectra were originally reduced with version 3.7 of the HARPS and HARPS-N Data Reduction Software, which includes color systematic corrections (Cosentino et al. 2014). RVs were computed using a numerical weighted mask following the methodology outlined by Baranne et al. (1996). Such a procedure has been shown to yield remarkable precision and accuracy (e.g., Mayor et al. 2009; Molaro et al. 2013; Pepe et al. 2013; Dumusque et al. 2014).

However, there is still room for improvement. A detailed analysis of the RV of individual spectral lines in HARPS spectra has revealed that some lines can show variations > 100 m s^{−1} as a function of time. These variations cannot be explained by stellar noise in a star like 55 Cnc. Indeed, stellar oscillations,

granulation phenomena, and stellar activity are expected to be of the order of a few m s^{−1} on an inactive G dwarf, as reported by Dumusque et al. (2011). After an in-depth study of the behavior of these lines, we identify three sources of velocity errors: the stitching of the CCD, some faint telluric lines, and fringing on the detector can all introduce significant velocity shifts of certain stellar lines that happen to fall near the stationary features, changing as the barycentric velocity for different observations scans the stellar lines across the artifacts.

The spectral lines affected by these variations were identified by calculating the periodogram of the RV of each spectral line used in the stellar template. All of the spectral lines that were exhibiting a signal more significant than 10% in false alarm probability were flagged as *bad* lines and removed from the stellar template. The result is a modified mask, which is used instead of the standard K5 mask to derive the RV of each observed spectrum by cross-correlation.

After performing a new cross-correlation of the data using this cleaned stellar template, the rms of the combined RV curve was reduced from 1.17 m s^{−1} to 1.04 m s^{−1}. In the rest of our analysis, we used the set of RV values from this revised reduction (a publication with this analysis is in preparation). Those RVs are presented in Table 1 and in Figure 1.

4. ESTIMATES OF LINE BROADENING DUE TO STELLAR ROTATION

The large number of high signal-to-noise ratio (S/N), high-resolution spectra gathered by HARPS-N allowed us to re-determine the stellar parameters of 55 Cnc using the Stellar Parameter Classification pipeline (Buchhave et al. 2014). We analyzed the 55 spectra observed on the night starting on 2013 November 14 with exposure times of 240 s and a resolution of $R = 115,000$, resulting in an average S/N per resolution element of 362 in the MgB region. We also analyzed three spectra, with S/N per resolution element of 143 and resolution $R = 48,000$, obtained between 2013 April 22 and 2014 March 8 with the fiber-fed Tillinghast Reflector Echelle Spectrograph (TRES; Szentgyorgyi & Furész 2007) on the 1.5 m Tillinghast Reflector at the Fred Lawrence Whipple Observatory on Mt. Hopkins, Arizona. The weighted average of each spectroscopic analysis yielded two sets of values from the two instruments, which are in very close agreement. The final stellar parameters are the average of these two sets, yielding $T_{\text{eff}} = 5358 \pm 50$ K, $\log(g) = 4.44 \pm 0.10$, and $[m/H] = 0.34 \pm 0.08$. Those values agree with the values published in the literature. The HARPS-N spectra yielded a projected rotational velocity of




Figure 1. Left: radial velocity data, in phase. HARPS data points are represented by inverted red triangles and HARPS-N by upright orange triangles. Right: the same data, binned in 14 equidistant points, and a model of the Rossiter–McLaughlin effect, for $v \sin i_* = 2.4 \text{ km s}^{-1}$ and $\beta = 0^\circ$.

(A color version of this figure is available in the online journal.)

Table 2
Analysis Priors and Results

Quantities (Units)	Winn et al. (2011)	Demory et al. (2011)	Nelson et al. (2014)	This Letter
Transit depth, ΔF (ppm)	380 ± 52	410 ± 63	...	384 ± 41
Impact parameter, b (R_*)	0.00 ± 0.24	0.16 ± 0.13	...	0.18 ± 0.08
Transit duration, W (days)	0.658 ± 0.0019	0.0665 ± 0.0019	...	0.0667 ± 0.0008
Mid-transit time, T_0 (BJD-2 450 000)	6184.50910 ± 0.00087	6184.5170 ± 0.0015	...	6184.5120 ± 0.0011
Period, P (days)	0.7365478 ± 0.0000014	0.7365478 ± 0.0000014
$\sqrt{V \sin i_*} \cos \beta$	-0.06 ± 0.42
$\sqrt{V \sin i_*} \sin \beta$	0.06 ± 0.69
Stellar $v \sin i_*$ (km s^{-1})	0.18 ± 0.48
Projected spin–orbit angle β (deg)	0 ± 180
Stellar rotation period, P_{rot} (d)	>20 (3σ)

Notes. The quantities determined by photometry are inserted as Gaussian priors taken from three papers. The values our analysis yields are included to illustrate that our fits did not force solutions to unrealistic values.

$v \sin i_* = 0.43 \text{ km s}^{-1}$ with an upper limit of 1.43 km s^{-1} , and the TRES spectra gave $v \sin i_* = 0.85 \text{ km s}^{-1}$ with an upper limit of 1.85 km s^{-1} . Both values agree with each other, but are lower than the value of $2.4 \pm 0.5 \text{ km s}^{-1}$ reported by Valenti & Fischer (2005).

We also computed the R'_{HK} activity index from our spectra and derived the age and rotation period of the star following Mamajek & Hillenbrand (2008). We arrive at an age for the star of $9.3 \pm 1.1 \text{ Gyr}$, a rotational period of $P_{\text{rot}} = 52 \pm 5 \text{ days}$, and a log of $R'_{\text{HK}} = -5.07 \pm 0.02$ using the HARPS-N spectra. The analysis of the HARPS spectra yields similar results, i.e., age = $10.0 \pm 1.2 \text{ Gyr}$, $P_{\text{rot}} = 54 \pm 5 \text{ days}$, and a log of $R'_{\text{HK}} = -5.11 \pm 0.02$. Both results agree with previously published values (von Braun et al. 2011; Dragomir et al. 2014) and imply a rotational velocity slower than 1 km s^{-1} , in agreement with our line broadening estimates.

5. ANALYSIS OF THE RADIAL VELOCITY DATA

The first step of our RV analysis consisted of removing the Keplerian orbital motion signals of the five planets discovered around 55 Cnc from each of our six data sets. We used the most recent orbital solution of the system obtained by Nelson et al. (2014). As recommended by these authors, we used the planetary masses and orbital parameters from their *Case 2*, in which they considered the errors of RV observations taken within 10 minutes of each other to be perfectly correlated. The corrected RVs for each data set are given in Table 1. Later tests using the Nelson et al. (2014) system parameters for their

Case 1, and also the planetary masses and orbital parameters of the system derived by Dawson & Fabrycky (2010) in their Tables 7, 8, and 10, yield similar results.

After removing the five planet signal, we modeled the Rossiter–McLaughlin effect for the six data sets combined using the formalism of Giménez (2006), based on Kopal (1942), and adjusted with a Monte Carlo Markov-Chain described in Triaud et al. (2013). All parameters determined by the photometry were controlled by priors issued from two independent data sets (see Table 2). Additional priors included the planet’s period (Nelson et al. 2014), the stellar parameters (von Braun et al. 2011), and the two $v \sin i_*$ values estimated in Section 4. We allowed the relative mean of each time series to float in order to absorb any offset produced by slightly different epochs of stellar activity (e.g., Triaud et al. 2009). Thirteen parameters were thus used on a total of 293 data points. A noise term of 70 cm s^{-1} was quadratically added to all measurement errors to reach a final reduced χ^2 of 0.97 ± 0.08 . All priors and important results are summarized in Table 2.

We conducted a number of different chains which all converge to the same conclusion: the Rossiter–McLaughlin effect is not detected (see Figure 1). The impossibility of constraining the spin–orbit angle of the system is illustrated in Figure 2, which shows the typical crescent shape expected when there is degeneracy between fast rotation and polar orbits and slow rotation and alignment (Triaud et al. 2011; Albrecht et al. 2011). This shape approximately maps contours of Rossiter–McLaughlin effects of equal semi-amplitudes. From the 3σ contour, we rule-out Rossiter–McLaughlin effects with semi-amplitudes larger




Figure 2. Posterior distribution for the projection of the stellar rotational speed, $v \sin i_*$, and the projection on the sky of the spin-orbit angle, β . 1, 2, and 3σ confidence contours are overplotted. Side histograms display the marginalized posteriors for each quantity. In the case of $v \sin i_*$, the two priors we applied are drawn as gray lines.

(A color version of this figure is available in the online journal.)

than 35 cm s^{-1} . The fact that anti-aligned orbits are as likely as aligned orbits implies that the Rossiter–McLaughlin effect is not detected.

The marginalized distribution in $v \sin i_*$ is thinner and peaks closer to 0 than our two estimates from spectral line broadening. Therefore, from the data we estimate $v \sin i_* = 0.2 \pm 0.5 \text{ km s}^{-1}$. This implies that the star’s most likely rotation period is 260 days (>22 days with 3σ confidence; >40 days only considering coplanar solutions).

To test the robustness of our analyses, we explored the impact of different priors. When replacing our two $v \sin i_*$ priors with the value estimated in Valenti & Fischer (2005), two symmetrical solutions, on polar orbits, are preferred: $\beta = +90^\circ$ and -90° (such a Rossiter–McLaughlin effect would have a semi-amplitude of 23 cm s^{-1}). A similar situation occurred for the spin-orbit angle measurement of WASP-80b (Triaud et al. 2013), which depends entirely on the value of $v \sin i_*$. Using no priors on $v \sin i_*$, the posterior is qualitatively similar, but both spikes are thinner and $v \sin i_*$ tails to higher values, as would be expected. The same procedures, removing the additional 70 cm s^{-1} noise added to the errorbars, produced similar results (with shorter confidence intervals).

6. DOPPLER TOMOGRAPHY

Given the impossibility of detecting the Rossiter–McLaughlin effect of 55 Cnc e, we tried to detect the signal of the planet using Doppler tomography (e.g., Albrecht et al. 2007; Collier Cameron et al. 2010). Doppler tomography reveals the distortion of the stellar line profiles when the planet, during transit, blocks part of the stellar photosphere. This distortion is a tiny dip in the stellar absorption profile, scaled down in width according

to the planet-to-star radius ratio. Additionally, the area of that dip corresponds to the planetary-to-stellar disks area ratio. As the planet moves across the stellar disk, the dip produces a trace in the time series of line profiles, which reveals the spin-orbit alignment between the star and planetary orbit.

For this analysis, we summed up all of the thousands of stellar absorption lines in each spectrum into one high S/N mean line profile. For this step, we employed a least-squares deconvolution (e.g. Collier Cameron et al. 2002) of the observed spectrum and theoretical lists of the stellar absorption lines from the Vienna Atomic Line Database (Kupka et al. 1999) for a star with $T_{\text{eff}} = 5200 \text{ K}$ and $\log g = 4.5$. The resulting line profiles were scaled so that their height was one, and were interpolated onto a velocity grid of 0.79 km s^{-1} increments, corresponding to the velocity range of one spectral pixel of HARPS-N at 550 nm. We then corrected the stellar line profiles for the RV of the host star and the barycentric velocity of the Earth. For each of the six runs, we summed up all the mean line profiles collected before and after the transit and subtracted the resulting profile from the in-transit ones. We then sorted the co-aligned line profiles of all the runs by orbital phase and combined the line profiles into one data set. Figure 3 shows the residuals of the line profiles and demonstrates that we are also unable to detect a trace of the transiting planet using this method. Our ability to detect the planet using Doppler tomography is in fact limited by the resolution of the spectra ($\sim 2.6 \text{ km s}^{-1}$ per resolution element), given the very slow rotational velocity of the star.

7. CONCLUSIONS

Our data do not support a detection of the spectroscopic transit of 55 Cnc e, but they rule out with 3σ confidence




Figure 3. Residuals of all line profiles of 55 Cnc taken during the six transits as a function of velocity and orbital phase of the planet. The two vertical dashed lines depict the area of the stellar line profile (FWHM). The units of the gray scale are fractional deviation from the average out-of-transit line profile. We are unable to detect the planetary signature in the line profiles.

any signal with a semi-amplitude larger than 35 cm s^{-1} . The non-detection can be explained by one of three scenarios: either 55 Cnc rotates more slowly than the average G-K main-sequence star ($\sim 2.4 \text{ km s}^{-1}$; Valenti & Fischer 2005), the orbital plane of the planets is perpendicular to the equatorial plane of the star, or the spin axis of the star is highly inclined with respect to us. Dragomir et al. (2014) report no variability associated with stellar spot rotation on a 42 day continuous monitoring of 55 Cnc with *MOST*. von Braun et al. (2011) estimated the system age at $10.2 \pm 2.5 \text{ Gyr}$. Therefore, the star is likely a very slow rotator, which is confirmed by our chromospheric activity measurements and stellar line broadening. Our conclusion is also consistent with a photometric modulation of 42.7 ± 2.5 days reported by Fischer et al. (2008), which they interpreted as stellar rotation. Because of the slow stellar rotation, reliably determining the projected spin-orbit angle of 55 Cnc e, or the inclination of its host’s spin axis, may remain out of reach. The slow stellar rotation also implies a weaker tidal coupling between 55 Cnc e and its host than what was presumed by Boué & Fabrycky (2014). Following Kaib et al. (2011), that would increase the probability of a spin-orbit misalignment.

While the detection of such a misalignment currently remains out of reach, our analysis confirms the capacity of the HARPS technology to reach a few tens of cm s^{-1} . These RV time series are the most constraining yet in precision and they reveal the suitability of HARPS and HARPS-N for follow-up and confirmation of small planets to be detected by missions like TESS. Such a level of RV precision will also be routinely reached by ESPRESSO on the Very Large Telescope. ESPRESSO will open up the study of weak Rossiter–McLaughlin effects, produced by super-Earths transiting slow rotators such as 55 Cnc e, with the potential to see if the same diversity that has been observed in the spin-orbit angle of hot Jupiters (Triaud et al. 2010; Brown et al. 2012; Albrecht et al. 2012) also exists for small planets.

Finally, Bourrier & Hébrard (2014) recently reported a detection of the Rossiter–McLaughlin effect of 55 Cnc e, with an amplitude of $\sim 60 \text{ cm s}^{-1}$. While such RV variations could be attributed to stellar surface physical phenomena (e.g., granulation or faculae), we argue that they are not produced by the Rossiter–McLaughlin effect, since the necessary $v \sin i_* \sim 3.3 \text{ km s}^{-1}$ is incompatible with the $v \sin i_*$, P_{rot} , age, and R'_{HK} activity index values we measure in Section 4. That $v \sin i_*$

is also incompatible with the $\sim 10 \text{ Gyr}$ stellar age derived by von Braun et al. (2011), and with the $P_{\text{rot}} > 40$ days rotational period estimates from Fischer et al. (2008) and Dragomir et al. (2014).

This publication was made possible through the support of a grant from the John Templeton Foundation. The opinions expressed are those of the authors and do not necessarily reflect the views of the John Templeton Foundation. The research leading to these results received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement 313014 (ETA-EARTH). This work used the VALD database, operated at Uppsala University, the Institute of Astronomy RAS in Moscow, and the University of Vienna. We thank the anonymous referee for a very constructive review and J. Winn for useful comments. A. H.M.J.T. is a Swiss National Science Foundation fellow under grant number P300P2-147773. F.R. acknowledges funding from the Alexander-von-Humboldt postdoctoral fellowship program. X.D. thanks the Swiss National Science Foundation (SNSF) for its support through an Early Postdoc Mobility fellowship. S.H. acknowledges support from the Spanish Ministry of Economy and Competitiveness under the 2011 Severo Ochoa Program MINECO SEV-2011-0187. HARPS-N is a collaboration between the Astronomical Observatory of the Geneva University, the CfA in Cambridge, the University of St. Andrews, the Queens University of Belfast, and the TNG-INAF Observatory. We thank all the researchers whose observing programs on HARPS were disrupted by our DDT program and who kindly observed for us. Finally, we thank ESO’s director general, Tim de Zeeuw, who granted our HARPS observing time.

Facilities: ESO:3.6m, TNG

REFERENCES

- Albrecht, S., Reffert, S., Snellen, I., Quirrenbach, A., & Mitchell, D. S. 2007, *A&A*, 474, 565
 Albrecht, S., Winn, J. N., Johnson, J. A., et al. 2011, *ApJ*, 738, 50
 Albrecht, S., Winn, J. N., Johnson, J. A., et al. 2012, *ApJ*, 757, 18
 Baranne, A., Queloz, D., Mayor, M., et al. 1996, *A&AS*, 119, 373
 Batygin, K., Morbidelli, A., & Tsiganis, K. 2011, *A&A*, 533, A7
 Boué, G., & Fabrycky, D. C. 2014, *ApJ*, 789, 111
 Bourrier, V., & Hébrard, G. 2014, arXiv:1406.6813
 Brown, D. J. A., Cameron, A. C., Anderson, D. R., et al. 2012, *MNRAS*, 423, 1503

- Buchhave, L. A., Bizzarro, M., Latham, D. W., et al. 2014, *Natur*, **509**, 593
- Chaplin, W. J., Sanchis-Ojeda, R., Campante, T. L., et al. 2013, *ApJ*, **766**, 101
- Collier Cameron, A., Bruce, V. A., Miller, G. R. M., Triaud, A. H. M. J., & Queloz, D. 2010, *MNRAS*, **403**, 151
- Collier Cameron, A., Horne, K., Penny, A., & Leigh, C. 2002, *MNRAS*, **330**, 187
- Cosentino, R., Lovis, C., Pepe, F., et al. 2012, *Proc. SPIE*, **8446**, 84461V
- Cosentino, R., Lovis, C., Pepe, F., et al. 2014, *Proc. SPIE*, 91478C
- Dawson, R. I., & Fabrycky, D. C. 2010, *ApJ*, **722**, 937
- Demory, B.-O., Gillon, M., Deming, D., et al. 2011, *A&A*, **533**, A114
- Dragomir, D., Matthews, J. M., Winn, J. N., & Rowe, J. F. 2014, in IAU Sym. 293, Formation, Detection, and Characterization of Extrasolar Habitable Planets, ed. N. Haghighipour (Cambridge: Cambridge Univ. Press), 52
- Dumusque, X., Bonomo, A. S., Haywood, R. D., et al. 2014, *ApJ*, **789**, 154
- Dumusque, X., Santos, N. C., Udry, S., Lovis, C., & Bonfils, X. 2011, *A&A*, **527**, A82
- Fischer, D. A., Marcy, G. W., Butler, R. P., et al. 2008, *ApJ*, **675**, 790
- Gaudi, B. S., & Winn, J. N. 2007, *ApJ*, **655**, 550
- Gillon, M., Demory, B.-O., Benneke, B., et al. 2012, *A&A*, **539**, A28
- Giménez, A. 2006, *ApJ*, **650**, 408
- Hirano, T., Narita, N., Shporer, A., et al. 2011, *PASJ*, **63**, 531
- Innanen, K. A., Zheng, J. Q., Mikkola, S., & Valtonen, M. J. 1997, *AJ*, **113**, 1915
- Kaib, N. A., Raymond, S. N., & Duncan, M. J. 2011, *ApJL*, **742**, L24
- Kopal, Z. 1942, *PNAS*, **28**, 133
- Kupka, F., Piskunov, N., Ryabchikova, T. A., Stempels, H. C., & Weiss, W. W. 1999, *A&AS*, **138**, 119
- Mamajek, E. E., & Hillenbrand, L. A. 2008, *ApJ*, **687**, 1264
- Mayor, M., Pepe, F., Queloz, D., et al. 2003, *Msngr*, **114**, 20
- Mayor, M., Udry, S., Lovis, C., et al. 2009, *A&A*, **493**, 639
- McArthur, B. E., Endl, M., Cochran, W. D., et al. 2004, *ApJL*, **614**, L81
- McLaughlin, D. B. 1924, *ApJ*, **60**, 22
- Molaro, P., Monaco, L., Barbieri, M., & Zaggia, S. 2013, *Msngr*, **153**, 22
- Mugrauer, M., Neuhäuser, R., Mazeh, T., et al. 2006, *AN*, **327**, 321
- Nelson, B. E., Ford, E. B., Wright, J. T., et al. 2014, *MNRAS*, **441**, 442
- Pepe, F., Cameron, A. C., Latham, D. W., et al. 2013, *Natur*, **503**, 377
- Queloz, D., Eggenberger, A., Mayor, M., et al. 2000, *A&A*, **359**, L13
- Rossiter, R. A. 1924, *ApJ*, **60**, 15
- Szentgyorgyi, A. H., & Furész, G. 2007, *RMxAA*, **28**, 129
- Triaud, A. H. M. J., Anderson, D. R., Collier Cameron, A., et al. 2013, *A&A*, **551**, A80
- Triaud, A. H. M. J., Collier Cameron, A., Queloz, D., et al. 2010, *A&A*, **524**, A25
- Triaud, A. H. M. J., Queloz, D., Bouchy, F., et al. 2009, *A&A*, **506**, 377
- Triaud, A. H. M. J., Queloz, D., Hellier, C., et al. 2011, *A&A*, **531**, A24
- Valenti, J. A., & Fischer, D. A. 2005, *ApJS*, **159**, 141
- von Braun, K., Boyajian, T. S., ten Brummelaar, T. A., et al. 2011, *ApJ*, **740**, 49
- Winn, J. N., Johnson, J. A., Howard, A. W., et al. 2010, *ApJL*, **723**, L223
- Winn, J. N., Matthews, J. M., Dawson, R. I., et al. 2011, *ApJL*, **737**, L18