



MIT Open Access Articles

Palladium-Catalyzed Synthesis of N-Aryl Carbamates

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation	Vinogradova, Ekaterina V., Nathaniel H. Park, Brett P. Fors, and Stephen L. Buchwald. "Palladium-Catalyzed Synthesis of N -Aryl Carbamates." <i>Org. Lett.</i> 15, no. 6 (March 15, 2013): 1394–1397.
As Published	http://dx.doi.org/10.1021/ol400369n
Publisher	American Chemical Society (ACS)
Version	Author's final manuscript
Citable link	http://hdl.handle.net/1721.1/93908
Terms of Use	Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.


Published in final edited form as:

Org Lett. 2013 March 15; 15(6): 1394–1397. doi:10.1021/ol400369n.

Palladium-Catalyzed Synthesis of N-Aryl Carbamates

 Ekaterina V. Vinogradova, Nathaniel H. Park, Brett P. Fors, and Stephen L. Buchwald
 Massachusetts Institute of Technology, Department of Chemistry, Cambridge, MA 02139, USA

Abstract



An efficient synthesis of aryl carbamates was achieved by introducing alcohols into the reaction of palladium-catalyzed cross-coupling of ArX (X = Cl, OTf) with sodium cyanate. The use of aryl triflates as electrophilic components in this transformation allowed for an expanded substrate scope for direct synthesis of aryl isocyanates. This methodology provides direct access to major carbamate protecting groups, S-thiocarbamates, and diisocyanate precursors to polyurethane materials.

Carbamates are found both in biologically active compounds¹ and in polymers,² and are valuable protecting groups in organic synthesis.³ As a result, many methods are available for their synthesis,⁴ including the aminolysis of acid anhydrides⁵ and chloroformates,⁶ the coupling of amines with CO₂ and alkyl halides,⁷ and the reaction of alcohols with isocyanates. These are often generated *in situ* in the presence of alcohols via the Hofmann,⁸ Curtius,⁹ Lossen,¹⁰ and Schmidt¹¹ rearrangements or the reductive carbonylation of nitroaromatic compounds.¹² These methods, however, suffer from a limited substrate scope, harsh reaction conditions, multiple-step preparations or the lack of readily available starting materials.


The shortcomings of the existing methods have prompted the development of new approaches to synthesize these important building blocks. Transition-metal catalyzed reactions of the isocyanate anion with aryl electrophiles or nucleophiles have shown potential for accessing the corresponding aryl isocyanate intermediates in the carbamate synthesis. Previously, Tkatchenko showed that Ni(0) catalysts promote the cross coupling of simple aryl halides with potassium cyanate.¹³ While this was the first example of using an isocyanate anion in a cross-coupling reaction, the yields of the carbamate products were low to moderate. More recently, a copper-catalyzed reaction of boronic acids with the same cyanate source has been reported by Baghersad.¹⁴ While the method allowed for improved yields of the carbamates, the substrate scope was limited in both the alcohols as well as the aromatic nucleophiles. During the completion of this work a related copper-catalyzed method for the *in situ* generation of aryl isocyanates has been introduced by Ma, where aryl bromides and iodides were used as electrophiles.¹⁵ While the substrate scope with respect to

sbuchwal@mit.edu.

Supporting Information Available. Experimental procedures, characterizations and spectral data for all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

the aryl halide was broad, the authors reported that they were unable to prepare *N*-Boc substituted amines. Furthermore, the range of other alcohols utilized was narrow.

Recently, we reported that the palladium-catalyzed cross-coupling of aryl chlorides and triflates with sodium cyanate affords aryl isocyanates that can be trapped *in situ* with amines to produce the corresponding ureas in a one-pot process.¹⁶ We felt that this could be extended to the synthesis of aryl carbamates by using alcohols as the nucleophilic trapping agents [Eq. (1)]. We now describe a general one-pot method for the palladium-catalyzed synthesis of aryl carbamates - including six of the most important carbamate protecting groups - from aryl chlorides and triflates.



(1)

For the synthesis of *N*-aryl ureas we demonstrated that the addition of phenol as a nucleophile to form a phenyl carbamate intermediate broadened the substrate scope of the process. We thought that the use of other alcohols would lead to general conditions for the synthesis of *N*-aryl carbamates.¹⁶ On this basis we began our current work by examining the Pd-catalyzed cross-coupling of 2,4-dimethoxy-5-chloropyrimidine with sodium cyanate using methanol as the nucleophilic trapping agent. No product of direct C–O cross-coupling reactions (alkyl-aryl ether) was observed and the desired *N*-aryl-*O*-alkyl carbamate was obtained in 86% yield (Table 1, entry **1a**).

It was necessary to preheat Pd₂dba₃ with **L1** prior to their introduction to the reaction mixture^{16,17} and to use triethylamine as an additive¹⁶ in order to obtain a high-yielding process. Using longer chain primary alcohols did not affect the course of the reaction and the corresponding carbamates were obtained in 82 - 92% yields (Table 1, entries **1b**, **1c**). Alcohols bearing trialkylsilyl functional groups could be employed, providing access to Teocarbamates (Table 1, entries **1d**, **1e**).

Moreover, the carbamate products were obtained in good yields (79 - 89%) when secondary alcohols were used (Table 1, entries **1f** - **1i**), including more sterically demanding alcohols with branching at the *b*-positions, such as 2-adamantyl alcohol and *L*-menthol (Table 1, entries **1g**, **1h**). Phenols could also be employed in the reaction,¹⁶ however, the stability of the resulting products is significantly lower. Isolation of *N*-aryl-*O*-phenyl carbamates was generally problematic as these products underwent significant decomposition during chromatographic purification on silica gel. However, with 4-fluorophenol the more stable carbamate could be isolated in 90% yield (Table 1, entry **1k**).¹⁸

The above-described cross-coupling conditions could be applied to an array of aryl electrophiles. Aryl chlorides bearing an ester, an unprotected acid or a secondary amide (Table 1, entries **1b**, **1g** and **1k**) were successfully converted to the corresponding *N*-aryl-*O*-alkyl carbamates. In our work on urea formation we showed that transmetallation of the isocyanate anion to the Pd-X intermediate was the most difficult step of the catalytic cycle.¹⁶ Thus, aryl bromides were found to be less reactive than the corresponding aryl chlorides or triflates. Here, too, the same trend was observed, although we were able to couple an electron-deficient aryl bromide, 3,5-bis(trifluoromethyl)bromobenzene; the corresponding carbamate product was obtained in 93% yield (Table 1, entry **1l**).

Aryl chlorides with one or two bulky *ortho*-substituents were generally less reactive under our standard conditions, producing the corresponding aryl carbamates in lower yields. Believing transmetalation to be the difficult step of the catalytic cycle, we felt that switching to aryl triflates might ameliorate this situation.^{16,19} In fact, the coupling reaction of 2,6-dimethylphenyl trifluoromethanesulfonate in the presence of 2-phenoxyethanol provided the carbamate product in 74% yield, while no product was observed with the corresponding aryl chloride (Table 1, entry **1j**).

We next sought to apply our method to the synthesis of the widely used *tert*-butyl carbamate (Boc) functional group. While the first set of conditions allowed for the efficient carbamate synthesis using primary and secondary alcohols, the use of 1.2 – 5-fold excess of *tert*-butanol afforded the corresponding Boc-protected anilines only in low yields (<50%). This result is consistent with the significantly lower reactivity of isocyanates towards tertiary alcohols.²⁰ However, employing *tert*-butanol as the solvent afforded the desired *tert*-butylcarbamates in 60 – 94% yield (Table 2).

Additionally there was no need either to preheat Pd₂dba₃ and **L1** or to use triethylamine as an additive. These reaction conditions tolerated aldehyde and cyano groups as well as a variety of heteroaryl electrophiles (Table 2, entries **2c-2i**).

Although the conditions described above (Tables 1 and 2) provided a process that manifested a broad substrate scope with respect to both aryl electrophile and the alcohol nucleophile, they were not suitable for use with certain alcohols, such as benzyl alcohols or trichloroethanol, and in these cases reduced yields of the carbamate products were observed. Additionally, byproducts resulting from direct C–S cross-coupling (thioethers) rather than aryl carbamates were observed when we attempted to use thiols as substrates. Thus, we attempted to utilize a phenyl carbamate, as before, followed by displacement with the alcohol or thiol (Scheme 1a).¹⁶ Unfortunately, this approach did not lead to a general protocol.

We felt that a more reactive intermediate, that could still be generated efficiently *in situ*, was needed in order to extend the scope of these reactions. Aryl isocyanates would be the ideal reactive intermediates needed for these reactions.²¹ Unfortunately in our previous work their synthesis was of limited success for sterically hindered aryl chlorides and with many heteroaryl chlorides.¹⁶

In fact, switching from aryl chlorides to aryl triflates allowed access to a broader range of aryl isocyanates (Scheme 1b). These include substrates bearing a variety of *ortho*-substituents (methyl, phenyl and isopropyl groups; Table 3, entries **3b, 3e-3g**), as well as other less reactive aryl-based electrophiles¹² (Table 3, entries **3c, 3d**). Trichloroethanol and benzyl alcohol were used to produce Troc- and Cbz-carbamates in 75% – 84% yields (Table 3, entries **3a-3d**). Butanethiol could also be used to provide the corresponding S-thiocarbamate in 86% yield (Table 3, entry **3e**). Additionally, the two-step approach enables the use of nucleophiles bearing otherwise reactive functional groups, such as aryl bromides, which could interfere with the cross-coupling process in the one-step procedure (Table 3, entry **3d**), thus allowing for further functionalization of the initial carbamate products.

Aryl diisocyanates are key intermediates in the industrial production of polyurethanes and are usually produced by phosgenation of the corresponding dianilines.²² To examine the potential of accessing these building blocks using our methodology, 5-chloro-2-methylphenyl trifluoromethanesulfonate was subjected to the one-pot two-step procedure. The *in situ* generated aryl diisocyanate was reacted with benzyl alcohol to afford the bis-Cbz-protected dianiline in 85% yield (92%/each carbamate formed; Table 3, entry **3f**).

9-fluorenylmethyl carbamate (Fmoc) is a protecting group for amino acids widely used in peptide synthesis. It can be easily cleaved by weak bases, such as piperidine and triethylamine.²³ Therefore, it is unsurprising that our standard conditions that employed triethylamine led to deprotection of the resulting carbamate and the formation of anilines as the major products. To overcome this problem, tris(2-(2-methoxyethoxy)ethyl)amine (TDA) was used in lieu of triethylamine.²⁴ The use of this milder base did not affect the stability of the resulting carbamate products and the Fmoc-protected amines **3g** and **3h** were obtained in 83% and 67% yields (Table 3). TDA was also employed for the cross-coupling of certain aryl triflates, as it was found to improve the yields of the corresponding carbamate products (Table 3, entries **3b**, **3c**, **3e**).

The synthesis of allyl carbamates typically requires the use of allyl chloroformate or allyl alcohol as the solvent, both of which are particularly hazardous substances. Therefore, applying the developed methodology to the synthesis of allyl carbamates would be advantageous, since it requires only a slight excess of the nucleophile. Allyl-esters and carbamates are known to easily undergo palladium(0)-catalyzed deprotection²⁵ in the presence of nucleophiles. Formation of the palladium *p*-allyl complex from the allyl carbamate generates the deprotonated carbamic acid, which readily decarboxylates to give the free amine.²⁶ In order to circumvent the undesired *p*-allyl pathway, the reactive palladium(0) species needed to be removed from the reaction mixture prior to the addition of allyl alcohol. Filtration of the reaction mixture through a plug of celite or silica gel did not remove the palladium species completely, which lead to diminished yields of the allyl carbamates. We envisioned that addition of an iodobenzene to the reaction mixture would provide an operationally simple procedure to rapidly convert the Pd(0) present to Pd(II), thus minimizing cleavage of the allyl carbamates without significantly altering the reaction conditions.²⁷ Indeed, stirring the reaction mixture with 10 mol % of iodobenzene prior to the introduction of allyl alcohol provided carbamates **3i** and **3j** in 63% and 67% yields (Table 3) using only 2-fold excess of the allyl alcohol.

In summary, we have shown that using alcohols as nucleophiles in the palladium-catalyzed cross-coupling of aryl chlorides and triflates with sodium isocyanate provides a broad range of aryl carbamate products in good yields. The substrate scope of the direct isocyanate coupling producing aryl isocyanates *in situ* was extended by using aryl triflates instead of aryl chlorides, allowing for the use of alcohols and thiols not compatible with cross-coupling conditions.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


Acknowledgments

We thank the National Institutes of Health (GM58160) for financial support of this project. N.H.P. acknowledges a National Science Foundation Graduate Research Fellowship. This activity was supported in part by an educational donation provided by Amgen. We thank James Colombe (M.I.T.) for helpful discussions and the Pentelute group (M.I.T.) for help in obtaining HRMS spectra. MIT has patents on the ligand used in this paper from which S.L.B. receives royalty payments.

References


1. a Baldinger SL, Shore ET. *Formulary*. 1996; 31:1029. b Nicolai, E.; Curtet, S.; Sicsic, J.; Lezoualc'h; Fischmeister, R.; Langlois, M.; Maillet, M.; Launay, M. 2007. U.S. Patent 7238693c Li W, Li J, Wu Y, Wu J, Hotchandani R, Cunningham K, McFadyen I, Bard J, Morgan P, Schlerman F, Xu X, Tam S, Goldman SJ, Williams C, Sypek J, Mansour TS. *J. Med. Chem.* 2009; 52:1799. [PubMed: 19278250]

2. a Lee BS, Chun BC, Chung Y-C, Sul KI, Cho JW. *Macromolecules*. 2001; 34:6431. b Król P. *Prog. Mater Sci.* 2007; 52:915.
3. Kocienski, PJ. *Protecting Groups: Thieme foundations of organic chemistry series*. Enders, D.; Noyori, R.; Trost, BM., editors. Thieme; Stuttgart; New York: 1994. p. 192-209.
4. Chaturvedi D. *Tetrahedron*. 2012; 68:15–45. and references therein.
5. Perron V, Abbott S, Moreau N, Lee D, Penney C, Zacharie B. *Synthesis*. 2008:283.
6. Carpino LA. *Acc. Chem. Res.* 1987; 20:401.
7. Salvatore RN, Shin SI, Nagle AS, Jung KW. *J. Org. Chem.* 2001; 66:1035. [PubMed: 11430069]
8. Keillor JW, Huang X. *Org. Syntheses*. 2002; 78:234.
9. Smith PAS. *Org. React.* 1946; 3:337.
10. Dubé P, Fine Nathel NF, Vetelino M, Couturier M, Larrivée Aboussafy C, Pichette S, Jorgensen ML, Hardink M. *Org. Lett.* 2009; 11:5622. [PubMed: 19908883]
11. Wolff H. *Org. React.* 1946; 3:307.
12. Paul F. *Coord. Chem. Rev.* 2000; 203:269.
13. Tkatchenko, I.; Jaouhari, R.; Bonnet, M.; Dawkins, G.; Lecolier, S. 1986. 1988. FR2575467 U.S. Patent 4749806
14. Kianmehr E, Baghersad MH. *Adv. Synth. Catal.* 2011; 353:2599.
15. Yang X, Zhang Y, Ma D. *Adv. Synth. Catal.* 2012; 354:2443.
16. Vinogradova EV, Fors BP, Buchwald SL. *J. Am. Chem. Soc.* 2012; 134:11132. [PubMed: 22716197]
17. For other uses of pre-mixing or pre-heating the palladium-source and ligand to access the active catalyst species prior to introduction into the reaction mixture, see: Ueda S, Su M, Buchwald SL. *J. Am. Chem. Soc.* 2012; 134:700. [PubMed: 22126442] Kotecki BJ, Fernando DP, Haight AR, Lukin KA. *Org. Lett.* 2009; 11:947. [PubMed: 19178160]
18. The electronic nature of the aryl alcohol does affect the efficiency of the transformation. The reactivity of substituted phenols was evaluated in experiments with 4-chlorotoluene. Using more electron-rich 4-methoxyphenol did not significantly alter the carbamate yield. Both phenyl *p*-tolylcarbamate and 4-methoxyphenyl *p*-tolylcarbamate were obtained in >95% yield, determined by ¹H NMR spectroscopy. However, no carbamate products were observed in the reactions with extremely electron-poor phenols, such as 4-cyanophenol, pentafluorophenol and 4-nitrophenol.
19. Hicks JD, Hyde AM, Cuezva AM, Buchwald SL. *J. Am. Chem. Soc.* 2009; 131:16720. [PubMed: 19886610]
20. Davis TL, Farnum JM. *J. Am. Chem. Soc.* 1934; 56:883.
21. Isocyanates are versatile intermediates, which can be used in a number of processes. For other examples, showing the extent of the isocyanate reactions, see aza-Wittig cyclizations: Marsden SP, McGonagle AE, McKeever-Abbas B. *Org. Lett.* 2008; 10:2589. [PubMed: 18489173] Rh(III)-catalyzed amidations: Hesp KD, Bergman RG, Ellman JA. *J. Am. Chem. Soc.* 2011; 133:11430. [PubMed: 21714533] 1,3-dipolar cycloadditions: Holt J, Fiksdahl A. *J. Heterocyclic Chem.* 2007; 44:375.
22. Seymour RB, Kauffman GB. *J. Chem. Educ.* 1992; 69:909.
23. Fields, GB. *Methods in Molecular Biology*. Pennington, MW.; Dunn, BM., editors. Vol. 35. Humana Press Inc.; Totowa, NJ: 1994. p. 17-27.
24. TDA has been previously shown to facilitate the cross coupling of aryl chlorides with sodium cyanate: [12], supporting information.
25. Jellerichs BG, Kong J-R, Krische MJ. *J. Am. Chem. Soc.* 2003; 125:7758. [PubMed: 12822967]
26. As expected, the presence of palladium in the reaction medium resulted in the deprotection and the formation of the corresponding anilines, in spite of the relatively mild conditions employed for the second step of the sequence (15 h at 45 °C).
27. Palladium(0)-complexes bearing biarylphosphine supporting ligands are known to undergo oxidative addition with sp² carbon-halogen bonds at room temperature. Halogen = Cl, Br: Wolfe JP, Singer RA, Yang BH, Buchwald SL. *J. Am. Chem. Soc.* 1999; 121:9550.



Scheme 1.
Synthesis of N-aryl carbamates, involving a) *O*-phenyl carbamate or b) isocyanate intermediates.

Table 1

Synthesis of aryl carbamates.^a

^aReaction conditions (isolated yields, average of 2 runs): ArX (1 mmol), NaOCN (2 mmol), ROH (1.2 mmol), Pd₂dba₃ (x mol %), **L1** (y mol %), NEt₃ (0.25 mmol), toluene (2 mL). The Pd₂(dba)₃ and **L1** were preheated in toluene at 120 °C for 3 minutes.

^bPd₂dba₃ (0.75 mol %), **L1** (1.8 mol %)

^cROH (2 mmol), 130 °C


^dPd₂dba₃ (0.5 mol %), **L1** (1.2 mol %)

^eNEt₃ (1.2 mmol), toluene (6 mL)

^ftoluene (8 mL)

Pd_2dba_3 (1.5 mol %), **L1** (3.6 mol %), 90 °C.

Table 2

Synthesis of Boc-carbamates.^a

^fPd₂dba₃ (0.75 mol %), **L1** (1.8 mol %), 24 h.

^aReaction conditions (isolated yields, average of 2 runs): ArX (1 mmol), NaOCN (2 mmol), Pd₂dba₃ (x mol %), **L1** (y mol %), *t*BuOH (2 mL).


^bPd₂dba₃ (1.5 mol %), **L1** (3.6 mol %)

^c90 °C

^dPd₂dba₃ (1.5 mol %), **L1** (3.6 mol %)

^e130 °C

Table 3

Synthesis of aryl carbamates.^{a,b}

^aReaction conditions (isolated yields, average of 2 runs). Step 1: ArX (1 mmol), NaOCN (2 mmol), Pd₂dba₃ (x mol %), **L1** (y mol %), toluene (2 mL). The Pd₂(dba)₃ and **L1** were preheated in toluene at 120 °C for 3 minutes. Step 2: NuH (2 mmol), NEt₃ (0.1 mmol).

^bTroc - 2,2,2-trichloroethoxycarbonyl, Cbz - carboxybenzyl, Fmoc - fluorenylmethyloxycarbonyl, Alloc - allyloxycarbonyl

^cNR₃ = NEt₃ (25 mol %)

^dPd₂dba₃ (0.5 mol %), **L1** (1.2 mol %)

^eTd₂dba₃ (1 mol %), **L1** (2.4 mol %)

^fNR₃ = TDA (10 mol %)

^g130 °C

^hPd₂dba₃ (0.75 mol %), **L1** (1.8 mol %)

ⁱNaOCN (3 mmol), toluene (3 mL). Step 2: BnOH (3 mmol), NEt₃ (0.2 mmol).

^jNo NEt₃ was used in Step 2.

^k0.5 mmol scale.

¹Step 2: PhI (0.1 mmol), 30 min at rt, then AllylOH (2 mmol), NEt₃ (0.1 mmol).