

PSFC/JA-99-34

Studies of EDA H-Mode in Alcator C-Mod

M. Greenwald, R. Boivin, P. Bonoli, C. Fiore, J. Goetz,
R. Granetz, A. Hubbard, I. Hutchinson, J. Irby,
Y. Lin, E. Marmor, A. Mazurenko, D. Mossessian,
T. Sunn Pedersen, J. Rice, J. Snipes, G. Schilling¹,
G. Taylor¹, J. Terry, S. Wolfe, S. Wukitch

November 1999

Plasma Science and Fusion Center
Massachusetts Institute of Technology
Cambridge, MA 02139

¹Princeton Plasma Physics Lab, Princeton, NJ 08543.

Submitted to Plasma Physics and Controlled Fusion.

This work was supported in part by the U. S. Department of Energy Contract No. DE-FC02-99ER54512. Reproduction, translation, publication, use and disposal, in whole or in part by or for the United States government is permitted.

STUDIES OF EDA H-MODE IN ALCATOR C-MOD

Abstract

Studies of the Enhanced $D\alpha$ H-mode (EDA) have been extended to include ohmic plasmas. No clear difference in the EDA/ELMfree boundary or in other phenomenology are seen between ohmic and ICRF heated plasmas, suggesting that neither the effects of ion tails nor direct RF/edge plasma interaction play a role in EDA. Edge safety factor (q_{95}) is the principal variable which determines which regime a discharge will be in. When q_{95} is above 4.0 for standard shaped plasmas, the discharge is almost always EDA and when it is below 3.5, the plasma is almost always ELMfree. New edge diagnostics have allowed measurement of pedestal profiles with resolution on the order of 1 mm. Sudden changes in profile widths are not seen when the plasma makes a transition from EDA to ELMfree however the widths do vary with the same parameters that determine the EDA/ELMfree boundary. Strong edge density fluctuations are observed to accompany EDA and may be responsible for the change in particle transport which is observed. The fluctuations have a quasi-coherent component whose frequency varies inversely with the pedestal width as measured by a visible continuum diagnostic.

1. Introduction

Alcator C-Mod is a compact high-field tokamak (minor radius = 0.22 m, major radius = 0.68 m, toroidal field = 3-8 T). The plasmas are shaped to match a closed divertor with $\kappa = 1.5 - 1.8$ and average triangularity ~ 0.4 . Plasma currents are typically in the range 0.4 - 1.4 MA. ICRF is the only auxiliary heating source, though at the high current densities that prevail, ohmic heating is never negligible. The plasma first wall is made of molybdenum tiles coated by in-vessel boronization.

Investigations of H-mode plasmas in Alcator C-Mod have centered on the EDA (Enhanced $D\alpha$) regime. The potential advantages of this regime have been well documented [1,2] and include good energy confinement, moderate particle confinement

(no impurity accumulation) and no large ELMs. Recent work has emphasized the scaling of the EDA/ELMfree boundary and attempts to uncover the underlying physics of the regime. Scans of plasma current and toroidal field show that EDA predominates at higher values of edge safety factor with a transition in the range $q_{95} = 3.5 - 4.0$ [3]. Plasma shape has also been shown to be important, with ELMfree more likely for triangularity below $\delta = 0.35$. There are some indications that ELMfree predominates above $\delta = 0.55$ as well. The EDA/ELMfree boundary is very weakly affected by neutral density with EDA more likely at midplane pressures above 0.015 PA. (When comparing to neutral densities in other devices, one must remember that the plasma density in C-Mod is about an order of magnitude higher than in low field tokamaks.) In some cases EDA discharges become ELMfree when input power is stepped down, however statistical studies of fully developed discharges do not show any obvious dependence.





Figure 1.. Traces from typical ohmic H-mode. B_T was ramped down to lower the L/H threshold (at 0.83 sec), then ramped up to raise q and allow the transition to EDA (at 1 sec.)

2. EDA in Ohmic plasmas

The only auxiliary heating employed on C-Mod is ICRF. Previous experiments reported on EDA H-modes were all with strong ICRF heating including a regime similar in many ways to EDA observed on JET in ICRF heated plasmas[4]. Despite previously reported C-Mod observations indicating that EDA was likely not a specific ICRF effect, these facts, combined with the apparent lack of EDA in neutral beam heated plasmas, has led to speculation that the heating method is crucial for its occurrence. However, new work on ohmic H-modes demonstrates that EDA is easily obtainable without ICRF heating. Figure 1 shows a number of traces from an ohmic EDA H-Mode. In this discharge, the L/H transition takes place at about 0.83 sec and the transition to EDA occurs at 1.0 sec. The lack of EDA in previous experiments on ohmic H-mode is understandable. Since the H-mode power threshold increases with B_T , ohmic H-modes are easiest to obtain at low q where the heating power is maximized for a given field. It was shown subsequently that EDA is favored by higher q . In the recent experiments, the toroidal field was ramped up after the L/H transition, raising q and accessing the EDA mode. The H-modes were maintained via a strong hysteresis effect seen in the power threshold. (In these cases, the H-mode could be maintained to levels where $P/nBS \sim 0.01$ (where P is measured in MW, n in $10^{20}/m^3$, B in T and S in m^2 .) Similar results were obtained by ramping the plasma current down following an L/H transition. Figure 2a and 2b show comparisons of the EDA/ELMfree boundary for ohmic and RF H-modes in which q_{95} is plotted against the mid-plane neutral pressure and total input power respectively. EDA is observed over a slightly wider range of q values for the ohmic discharges though the difference is probably not significant. Considering a wider set of data than shown in the figure, there is evidence for a very weak dependence on neutral and plasma density. Despite anecdotal indications, no clear dependence on input power or net power across the separatrix is observed. The implication of obtaining EDA under the same conditions in ICRF and ohmic plasmas is that the physics associated with a particular heating scheme can be ruled out as a cause for EDA. These effects could have included the dynamics of energetic ion tails or the interaction between ICRF waves and the plasma edge. The former was never likely, as the high densities that prevail in C-Mod prevent ion tails of

significant energy from forming. Earlier work which found a dependence on triangularity could not be tested in this study, since there was no variation in plasma shape for the ohmic H-modes.



2a. A comparison of the EDA/ELMfree boundary for ohmic (filled symbols) and ICRF heated discharges (open symbols). There is no significant difference between the two. With this data set, no dependence on midplane neutral pressure can be seen, though some small effect is seen when a larger data set is considered.

2b. A comparison of the EDA/ELMfree boundary plotted against input power. Ohmic and ICRF heated plasmas again show the same dependence on q_{95} . No obvious trend with input power is apparent.

3. Pedestal measurements

Because of its compact size, high field and high density, length scales - major radius, gyroradius, connection length, mean free path - are all smaller in C-Mod than in other devices. It therefore came as no surprise that pedestal widths in C-Mod H-modes were

smaller as well, initially challenging the instrumental resolution of the edge diagnostics[5]. During the past two years, a number of diagnostics with significantly improved spatial resolution has been installed. These include an edge Thomson scattering system with resolution of approximately 1.3 mm mapped to the midplane; a visible bremsstrahlung (VB) array with 0.6 mm resolution; a pair of soft x-ray arrays with 1.5 mm resolution; and a Ly α array, which can measure the ionization rate, which has 2 mm resolution. Further details on these new diagnostics can be found in reference [6]. An ECE polychromator is also available. While its resolution is no better than about 8 mm, it seems sufficient to resolve the edge electron temperature profile in H-mode which are in the range 6-12 mm. The pedestal width for electron density, on the other hand, seems to be much narrower - typically in the range of 2-4 mm. The VB array which observes continuum radiation whose emission scales like $n^2 Z_{\text{EFF}}/g(\text{Te})$ (and thus is most sensitive to density profiles) also sees narrow pedestal widths, but with greater variation, ranging from 2-15 mm. The soft x-ray emission is believed to come from recombination of fully stripped fluorine and shows widths from 2-9 mm. Relatively little variation is seen in the electron temperature width; in particular, no difference is seen between temperature profile widths from EDA and ELMfree H-modes. The most significant dependence observed is an increase in Te pedestal width as the net power through the separatrix increases[5]. No sharp difference is seen in the pedestal widths of any of these diagnostics as the plasma makes transitions between EDA and ELMfree H-mode. All, however, show some variation in direct proportion to δ and q_{95} , the variables which determine the EDA/ELMfree boundary. The VB and x-ray widths generally show the greatest increases. Figure 3 shows traces of H α and the pedestal widths for a shot in which triangularity was scanned, causing the discharge to make a transition from EDA to ELMfree. On the same graph, one can see that there is less change in the density profile width as measured by Thomson scattering. The cause of this difference is not yet understood.

4. Fluctuations

In EDA, the edge pressure gradients can be at or above the ideal ballooning limit but are not relaxed by type I ELMs; instead a continuous process would seem to be at work. This process is probably related to broadband and quasi-coherent fluctuations which are




Figure 3.. Pedestal widths from various diagnostics are plotted for a discharge where the triangularity was scanned from 0.6 to 0.28. Because of low input power, the H-mode was interrupted by back transitions to L-mode and periods of type III ELMs. (No data is plotted during these intervals.) None the less, the widths show a clear co-variance with triangularity.

observed during EDA. These fluctuations are seen with reflectometry, PCI (Phase Contrast Imaging) and magnetic pick-up loops. The coherent component has a frequency on the order of 100 kHz with significant variation during and between shots. In general higher frequencies are seen as the plasma transitions into or out of EDA. Plasmas with some ELMfree characteristics (longer impurity confinement times) have higher frequency coherent oscillations than those which are most clearly EDA. The reflectometer localizes the oscillations to the plasma edge where the Doppler shift in the shear layer is likely to make an important contribution to the lab frame frequency. The




Figure. 4. The frequency of the quasi-coherent fluctuation observed by reflectometry is plotted along with the inverse of the pedestal width as observed by visible bremsstrahlung radiation. The dependence on inverse scale length suggests a diamagnetic origin for the fluctuation frequency.

fluctuation frequency is seen to vary inversely with the pedestal scale length as measured by VB (fig 4) suggesting a diamagnetic origin. The coherent feature is dominant in the PCI measurements, which show a well defined spatial structure with $k_R \sim 6 \text{ cm}^{-1}$. PCI makes measurements averaged through a vertical chord, so with the observed fluctuations localized in the plasma edge by reflectometry, the measured k_R corresponds to k_θ . Magnetic diagnostics typically observe only the broadband fluctuations suggesting that the coherent feature either has a short wavelength ($< 1\text{-}3 \text{ cm}$) consistent with PCI measurements, or has no significant magnetic component. In a few cases, weakly coherent fluctuations in the same frequency band as those discussed above, are seen on pickup coils located near the inner edge (small major radius) of the plasma. Coils on the outside never see this component of the fluctuations, possibly because they are 2 cm farther from the plasma.

5. Discussion

EDA could be an interesting mode for a fusion reactor. It avoids the twin perils of the H-mode; impurity accumulation when ELMfree and high pulsed heat loads on the divertor

in ELMy regimes. However, we cannot extrapolate the mode to a reactor with any confidence unless we understand the underlying physics. A number of effects seems to be ruled out. As discussed above, specific ICRF physics appears to play no role in the phenomenon. The lack of any important dependence on neutral pressure or density suggest that neutral losses or viscosity are unimportant. The strong dependences observed on q_{95} and triangularity [3] on the other hand, would seem to implicate MHD stability in the process. So far, however, calculations of ballooning stability find no clear trends that align with the EDA/ELMfree boundary. Moreover, when radiation or neutral pressures are high, low performance EDA discharges are obtained. These have very poor confinement and small pedestals and are apparently quite far from the stability limits. It must be noted that these calculations are incomplete since current density profiles in the edge plasma are not yet available.

An interesting subsidiary question is whether the EDA represents a separate stable regime which is attained through a bifurcation process or whether it can be joined continuously to the ELMfree regime. Evidence on this point are inconclusive. On some discharges, the transition is accompanied by an abrupt change in $H\alpha$; on others it is not. Pedestal measurements do not, in general, show any abrupt change. On the other hand, impurity confinement does change dramatically and the density fluctuations, discussed above, turn on abruptly[3]. Dithers between EDA and ELMfree regimes, some lasting for as short a time as 1-2 msec, are seen on $H\alpha$ and on the density fluctuation diagnostics suggesting limit cycle behavior.

Recent simulations of drift Alfvén turbulence [7] show a coherent mode grow up as the pedestal pressure gradient approaches an MHD stability limit which can be several times the ideal limit. The predicted mode has $k_{\theta} \sim 2\pi/L_P$ where L_P is the pedestal gradient length, consistent with the C-Mod observations. The frequency predicted is of the correct magnitude and dependence. The critical tests for this identification are: 1. does the mode preferentially drive particle transport? and 2. are the dependences on shape and q reproduced? Answers to these questions will require further analysis and addition of more realistic geometry to the simulations.

Acknowledgements

The authors wish to thank the entire Alcator staff for their hard work and dedication in support of these experiments. Work supported by U.S. Department of Energy contract DE-FC02-99ER54512.

References

- [1]. J. Snipes, et al., 24th EPS – Berchtesgaden, Vol. 21A, Part II, 565 (1997)
- [2]. M. Greenwald, et al., Nucl. Fusion **37**, 793 (1997)
- [3]. M. Greenwald, et al., Phys. Plasmas, **6**, 1943 (1999)
- [4]. M. Bures, et al, Nucl. Fus., **32**, 539 (1992)
- [5]. A. Hubbard, et al., to be published in the proceedings of the 26th conference on Plasma Physics and Controlled Fusion in Maastricht, (1999)
- [6]. D. Mossessian, et al., Proceedings of this conference, submitted to Plasma Phys. and Cont. Fusion.
- [7]. B. Rogers, J. Drake, Phys. Plasmas, **6**, 2797 (1999) (1999)