

MIT Open Access Articles

Editorial: Cell and molecular signaling, and transport pathways involved in growth factor control of synaptic development and function

The MIT Faculty has made this article openly available. ***Please share*** how this access benefits you. Your story matters.

Citation: Yoshii, Akira, Martha Constantine-Paton, and Nancy Y. Ip. "Editorial: Cell and Molecular Signaling, and Transport Pathways Involved in Growth Factor Control of Synaptic Development and Function." *Frontiers in Synaptic Neuroscience* 7 (June 4, 2015).

As Published: <http://dx.doi.org/10.3389/fnsyn.2015.00008>

Publisher: Frontiers Research Foundation

Persistent URL: <http://hdl.handle.net/1721.1/98181>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of use: Creative Commons Attribution



Editorial: Cell and molecular signaling, and transport pathways involved in growth factor control of synaptic development and function

Akira Yoshii^{1*}, Martha Constantine-Paton^{2,3,4} and Nancy Y. Ip^{5,6,7}

¹ Department of Anatomy and Cell Biology, and Pediatrics, University of Illinois at Chicago, Chicago, IL, USA, ² Department of Brain and Cognitive Science, McGovern Institute for Brain Research, Massachusetts Institute of Technology, Cambridge, MA, USA, ³ McGovern Institute for Brain Research and Massachusetts Institute of Technology, Cambridge, MA, USA,

⁴ Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, USA, ⁵ Division of Life Science, Hong Kong University of Science and Technology, Hong Kong, China, ⁶ Molecular Neuroscience Center and Hong Kong University of Science and Technology, Hong Kong, China, ⁷ State Key Laboratory of Molecular Neuroscience, Hong Kong University of Science and Technology, Hong Kong, China

Keywords: BDNF, neurotrophic factors, synapse formation, FGF7, FGF22, UNC-5, UNC-6 Netrin

OPEN ACCESS

Edited by:

Per Jesper Sjöström,
The Research Institute of the McGill
University Health Centre, Canada

Reviewed by:

Keith Murai,
McGill University, Canada

*Correspondence:

Akira Yoshii,
ayoshii@uic.edu

Received: 08 May 2015

Accepted: 20 May 2015

Published: 04 June 2015

Citation:

Yoshii A, Constantine-Paton M and Ip NY (2015) Editorial: Cell and molecular signaling, and transport pathways involved in growth factor control of synaptic development and function. *Front. Synaptic Neurosci.* 7:8. doi: 10.3389/fnsyn.2015.00008

Since the discovery of nerve growth factor (NGF) more than a half century ago (Levi-Montalcini and Cohen, 1960), the prototypic neurotrophin family has included brain derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3), and neurotrophin-4 (NT-4). Neurotrophins bind to the Trk family of receptors, as well as the p75 receptor, to activate multiple intracellular signaling cascades (reviewed by Reichardt, 2006). BDNF receptor tropomyosin receptor kinase B (TrkB) signaling has been extensively studied for its roles in the central nervous system (CNS) ranging from cell survival, axonal and dendritic growth and synapse formation. The pathway mediates long-lasting activity-modulated synaptic changes on excitatory and inhibitory neurons and plays critical roles in circuit development and maintenance. In addition to BDNF, many studies have identified other “growth” or signaling factors in the CNS that play important roles in the development, maintenance, and control of synaptic and circuit function. However, details of the intracellular signaling systems downstream of these events are frequently unexplored. In this Research Topic, we have collected original studies and review articles that present cellular and molecular mechanisms concerning activity-dependent synapse formation and their implications for behavior and brain disorders.

Vadodaria and Jessberger discuss synapse maturation in adult-born dentate granule cells and the role of BDNF-TrkB and several other signaling pathways that activate Cdc42, Rac1, and RhoA (Vadodaria and Jessberger, 2013). These small Rho GTPases regulate polymerization of actin and microtubules, and are consequently involved in aspects of neuronal maturation ranging from cell migration, to dendritic arborization, spine maturation, and synaptic integration of these newborn hippocampal neurons.

Kellner et al. show that BDNF is critical for activity-dependent maintenance of mature spines through F-actin polymerization (Kellner et al., 2014). Integrity of this spine cytoskeleton is also critical for the vesicular transport, carried out by molecular motor proteins. For example, Myosin Va is a plus end actin, vesicular motor protein that carries postsynaptic density protein 95 (PSD-95), Synapse-associated protein 90/postsynaptic density-95-associated protein (SAPAP) and Shank, an essential glutamate receptor scaffold complex along actin to the postsynaptic membrane at the tip of dendritic spines (Hammer and Wagner, 2013; Yoshii et al., 2013). Furthermore, BDNF-TrkB signaling triggers transport of the vesicles containing PSD-95 and its associated complex

to postsynaptic membranes via activation of PI3-kinase-Akt pathway (Yoshii and Constantine-Paton, 2007).

BDNF-TrkB signaling pathway also regulate the molecular assembly of synaptic membrane. Zonta and Minichiello discuss the emerging role of lipid raft, the detergent resistant lipid microdomain enriched with cholesterol and sphingolipid, in synaptic plasticity as a result of neurotrophin signaling (Zonta and Minichiello, 2013). When BDNF binds to the TrkB receptor, the ligand-receptor complex shifts to lipid rafts via activation of tyrosine kinase Fyn (Pereira and Chao, 2007; Suzuki et al., 2007). Lipid raft is enriched in postsynaptic membrane, and facilitates localization of palmitoylated proteins such as PSD-95, which is a critical regulator of synaptic plasticity at excitatory synapses. Palmitoylation of PSD-95 in the cell body is also mediated by BDNF-TrkB signaling via activation of phospholipase C γ (PLC γ) and protein kinase C (PKC). The PKC inhibitors chelerythrine as well as a synthetic zeta inhibitory peptide (ZIP) designed to block the brain-specific PKC isoform protein kinase M ζ (PKM ζ) were used to demonstrate this effect (Yoshii et al., 2011). However, additional studies in the hippocampus began to raise concerns about the specificity of ZIP (Lee et al., 2013; Volk et al., 2013). A follow-up study by Yoshii and Constantine-Paton in this Topic confirmed that, while both chelerythrine and ZIP could suppress the postsynaptic localization of PSD-95, PKM ζ knock-down with RNA interference did not exhibit this effect. The result indicates that the ZIP peptide, widely used as a “specific” PKM ζ antagonist, may block another PKC variant that is the kinase actually involved in PSD-95 palmitoylation in cell body.

BDNF is critical for mechanisms underlying various modalities of sensory processing, cognition and behaviors. These roles have been studied in hippocampal learning and memory (reviewed by Minichiello, 2009), in maturation and plasticity of the CNS and also in neurological disease (reviewed by Yoshii and Constantine-Paton, 2010). It is not surprising that BDNF is associated with brain disorders such as epilepsy, autism, depression, and schizophrenia since all of these have chronic effects on synaptic function. However, Andersen and Sonntag have studied the effect of juvenile exposure to psychostimulants on the risk of cocaine addiction in adulthood. They found that treatment with methylphenidate, frequently used to improve attention in children, has a long-lasting suppressive effect on cocaine-induced increases in BDNF expression (Andersen and Sonntag, 2014). Further studies will hopefully identify the mechanisms underlying the critical period effect on addiction as well as long term effect of stimulants.

Neurotrophic factors also affect feeding. Maekawa et al. have shown that low BDNF expression in the ventromedial hypothalamus correlates with blood glucose level, increased leptin secretion and eating, and visceral fat accumulation in a type 2 diabetic rat line (Maekawa et al., 2013). The results indicate BDNF and leptin play major roles in central regulation of energy metabolism and dysregulation of the neurotrophin signaling result in obesity.

Harvey discusses that leptin regulates synaptic functions (Harvey, 2013). This dietary hormone activates extracellular-signal-regulated kinase (ERK) and facilitates GluN2B-mediated synaptic depression during early postnatal development while it regulates LTP through PI3-kinase pathway in adult hippocampus (Moult and Harvey, 2011).

It is now clear that variety of synaptogenic growth factors are wider than previously thought. Two articles review diversity of these factors. Williams and Umemori discuss members of the fibroblast growth factor (FGF) family in the context of synaptic development (Williams and Umemori, 2014). FGFs have been shown to organize presynaptic vesicle clustering. Remarkably, FGF7-null mice exhibit a specific deficit in hippocampal inhibitory synapse formation while FGF22-null mice are deficient in excitatory synapses (Terauchi et al., 2010). Furthermore, FGF7-null animals are prone to develop epilepsy after kindling, while FGF22-null are resistant to this seizure induction.

Poon et al. provide a comprehensive review of Netrin, Wnt, transforming growth factor- β (TGF- β), tumor necrosis factor- α (TNF- α), all of which were first identified for their roles other than synaptogenesis (Poon et al., 2013). For example, UNC-6/netrin and its receptor UNC-5 were originally described as axonal guidance molecules. However, UNC-6 and UNC-5 facilitate localization of presynaptic proteins to axons by excluding them from dendrites (Poon et al., 2008). Consequently, these pathway are all involved in regulating axonodendritic polarity.

The field of neuronal growth factors is continuing to grow, and new discoveries, some which are highlighted in this volume, will prompt new questions. For example, do these growth factors work together, competitively or separately? Is there cell-type specificity for each factor? Do they play a deterministic or a modulatory role in synaptic specificity during development? Advances in genomics and proteomics will help us understand not only single cascade but also multiple signaling pathways as network. Various genetic tools allow spacial and temporal controls of gene expression and neuronal activities (Boyden et al., 2005; Arenkiel and Ehlers, 2009; Konermann et al., 2013). Super-resolution microscopy enables observations of signaling molecules at synapse and organelles in unprecedented manner. Multi-photon microscopy has been invaluable to study a wide range of structures from individual spines to neuronal circuitry. Applications of these new technologies will create exciting opportunities to tackle the above and other questions.

Acknowledgments

This work was supported by a DoD grant W81XWH-09-1-0088 to AY, NIH grants 5R01EY006039-27 and 5R01EY014074-15 to MC-P, and the National Key Basic Research Program of China (2013CB530900), Hong Kong Research Grants Council Theme-based Research Scheme (T13-607/12R), and Shenzhen Peacock Plan to NYI.

References

- Andersen, S. L., and Sonntag, K. C. (2014). Juvenile methylphenidate reduces prefrontal cortex plasticity via D3 receptor and BDNF in adulthood. *Front. Synaptic Neurosci.* 6:1. doi: 10.3389/fnsyn.2014.00001
- Arenkiel, B. R., and Ehlers, M. D. (2009). Molecular genetics and imaging technologies for circuit-based neuroanatomy. *Nature* 461, 900–907. doi: 10.1038/nature08536
- Boydén, E. S., Zhang, F., Bamberg, E., Nagel, G., and Deisseroth, K. (2005). Millisecond-timescale, genetically targeted optical control of neural activity. *Nat. Neurosci.* 8, 1263–1268. doi: 10.1038/nn1525
- Hammer, J. A., and Wagner, W. (2013). Functions of class V myosins in neurons. *J. Biol. Chem.* 288, 28428–28434. doi: 10.1074/jbc.R113.514497
- Harvey, J. (2013). Leptin regulation of neuronal morphology and hippocampal synaptic function. *Front. Synaptic Neurosci.* 5:3. doi: 10.3389/fnsyn.2013.00003
- Kellner, Y., Gödecke, N., Dierkes, T., Thieme, N., Zagrebelsky, M., and Korte, M. (2014). The BDNF effects on dendritic spines of mature hippocampal neurons depend on neuronal activity. *Front. Synaptic Neurosci.* 6:5. doi: 10.3389/fnsyn.2014.00005
- Konermann, S., Brigham, M. D., Trevino, A. E., Hsu, P. D., Heidenreich, M., Cong, L., et al. (2013). Optical control of mammalian endogenous transcription and epigenetic states. *Nature* 500, 472–476. doi: 10.1038/nature12466
- Lee, A. M., Kanter, B. R., Wang, D., Lim, J. P., Zou, M. E., Qiu, C., et al. (2013). Prkcz null mice show normal learning and memory. *Nature* 493, 416–419. doi: 10.1038/nature11803
- Levi-Montalcini, R., and Cohen, S. (1960). Effects of the extract of the mouse submaxillary salivary glands on the sympathetic system of mammals. *Ann. N.Y. Acad. Sci.* 85, 324–341.
- Maekawa, F., Fujiwara, K., Toriya, M., Maejima, Y., Nishio, T., Toyoda, Y., et al. (2013). Brain-derived neurotrophic factor in VMH as the causal factor for and therapeutic tool to treat visceral adiposity and hyperleptinemia in type 2 diabetic Goto-Kakizaki rats. *Front. Synaptic Neurosci.* 5:7. doi: 10.3389/fnsyn.2013.00007
- Minichiello, L. (2009). TrkB signalling pathways in LTP and learning. *Nat. Rev. Neurosci.* 10, 850–860. doi: 10.1038/nrn2738
- Moult, P. R., and Harvey, J. (2011). NMDA receptor subunit composition determines the polarity of leptin-induced synaptic plasticity. *Neuropharmacology* 61, 924–936. doi: 10.1016/j.neuropharm.2011.06.021
- Pereira, D. B., and Chao, M. V. (2007). The tyrosine kinase Fyn determines the localization of TrkB receptors in lipid rafts. *J. Neurosci.* 27, 4859–4869. doi: 10.1523/JNEUROSCI.4587-06.2007
- Poon, V. Y., Choi, S., and Park, M. (2013). Growth factors in synaptic function. *Front. Synaptic Neurosci.* 5:6. doi: 10.3389/fnsyn.2013.00006
- Poon, V. Y., Klassen, M. P., and Shen, K. (2008). UNC-6/netrin and its receptor UNC-5 locally exclude presynaptic components from dendrites. *Nature* 455, 669–673. doi: 10.1038/nature07291
- Reichardt, L. F. (2006). Neurotrophin-regulated signalling pathways. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 361, 1545–1564. doi: 10.1098/rstb.2006.1894
- Suzuki, S., Kiyosue, K., Hazama, S., Ogura, A., Kashihara, M., Hara, T., et al. (2007). Brain-derived neurotrophic factor regulates cholesterol metabolism for synapse development. *J. Neurosci.* 27, 6417–6427. doi: 10.1523/JNEUROSCI.0690-07.2007
- Terauchi, A., Johnson-Venkatesh, E. M., Toth, A. B., Javed, D., Sutton, M. A., and Umemori, H. (2010). Distinct FGFs promote differentiation of excitatory and inhibitory synapses. *Nature* 465, 783–787. doi: 10.1038/nature09041
- Vadodaria, K. C., and Jessberger, S. (2013). Maturation and integration of adult born hippocampal neurons: signal convergence onto small Rho GTPases. *Front. Synaptic Neurosci.* 5:4. doi: 10.3389/fnsyn.2013.00004
- Volk, L. J., Bachman, J. L., Johnson, R., Yu, Y., and Huganir, R. L. (2013). PKM- ζ is not required for hippocampal synaptic plasticity, learning and memory. *Nature* 493, 420–423. doi: 10.1038/nature11802
- Williams, A. J., and Umemori, H. (2014). The best-laid plans go oft awry: synaptogenic growth factor signaling in neuropsychiatric disease. *Front. Synaptic Neurosci.* 6:4. doi: 10.3389/fnsyn.2014.00004
- Yoshii, A., and Constantine-Paton, M. (2007). BDNF induces transport of PSD-95 to dendrites through PI3K-AKT signaling after NMDA receptor activation. *Nat. Neurosci.* 10, 702–711. doi: 10.1038/nn1903
- Yoshii, A., and Constantine-Paton, M. (2010). Postsynaptic BDNF-TrkB signaling in synapse maturation, plasticity, and disease. *Dev. Neurobiol.* 70, 304–322. doi: 10.1002/dneu.20765
- Yoshii, A., Murata, Y., Kim, J., Zhang, C., Shokat, K. M., and Constantine-Paton, M. (2011). TrkB and protein kinase M ζ regulate synaptic localization of PSD-95 in developing cortex. *J. Neurosci.* 31, 11894–11904. doi: 10.1523/JNEUROSCI.2190-11.2011
- Yoshii, A., Zhao, J. P., Pandian, S., van Zundert, B., and Constantine-Paton, M. (2013). A myosin va mutant mouse with disruptions in glutamate synaptic development and mature plasticity in visual cortex. *J. Neurosci.* 33, 8472–8482. doi: 10.1523/JNEUROSCI.4585-12.2013
- Zonta, B., and Minichiello, L. (2013). Synaptic membrane rafts: traffic lights for local neurotrophin signaling? *Front. Synaptic Neurosci.* 5:9. doi: 10.3389/fnsyn.2013.00009

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Yoshii, Constantine-Paton and Ip. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.