

MIT Open Access Articles

Joint effects of storm surge and sea-level rise on US Coasts: new economic estimates of impacts, adaptation, and benefits of mitigation policy

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Neumann, James E., Kerry Emanuel, Sai Ravela, Lindsay Ludwig, Paul Kirshen, Kirk Bosma, and Jeremy Martinich. "Joint Effects of Storm Surge and Sea-Level Rise on US Coasts: New Economic Estimates of Impacts, Adaptation, and Benefits of Mitigation Policy." *Climatic Change* 129, no. 1–2 (December 14, 2014): 337–349.

As Published: <http://dx.doi.org/10.1007/s10584-014-1304-z>

Publisher: Springer-Verlag

Persistent URL: <http://hdl.handle.net/1721.1/99157>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-Noncommercial-Share Alike



Climatic Change

Joint Effects of Storm Surge and Sea-level Rise on US Coasts

--Manuscript Draft--

Manuscript Number:	
Full Title:	Joint Effects of Storm Surge and Sea-level Rise on US Coasts
Article Type:	Research Article
Corresponding Author:	James Neumann Industrial Economics, Incorporated Cambridge, MA UNITED STATES
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Industrial Economics, Incorporated
Corresponding Author's Secondary Institution:	
First Author:	James Neumann
First Author Secondary Information:	
Order of Authors:	James Neumann Kerry Emanuel Sai Ravela Lindsay Ludwig Paul Kirshen Kirk Bosma Jeremy Martinich
Order of Authors Secondary Information:	
Abstract:	Recent literature, the US Global Change Research Program's National Climate Assessment, and recent events, such as Hurricane Sandy, highlight the need to take better account of both storm surge and sea-level rise (SLR) in assessing coastal risks of climate change. This study combines three models - a tropical cyclone simulation model; a storm surge model; and a model for economic impact and adaptation - to estimate the joint effects of storm surge and SLR for the US coast through 2100. The model is tested using multiple SLR scenarios, including those incorporating estimates of dynamic ice-sheet melting, two global greenhouse gas (GHG) mitigation policy scenarios, and multiple general circulation model climate sensitivities. The results illustrate that a large area of coastal land and property is at risk of damage from storm surge today; that land area and economic value at risk expands over time as seas rise and as storms become more intense; that adaptation is a cost-effective response to this risk, but residual impacts remain after adaptation measures are in place; that incorporating site-specific episodic storm surge increases national damage estimates by a factor of two relative to SLR-only estimates, with greater impact on the East and Gulf coasts; and that mitigation of GHGs contributes to significant lessening of damages. For a mid-range climate-sensitivity scenario that incorporates dynamic ice sheet melting, the approach yields national estimates of the impacts of storm surge and SLR of \$990 billion through 2100 (net of adaptation, cumulative undiscounted 2005\$); GHG mitigation policy reduces the impacts of the mid-range climate-sensitivity estimates by \$84 to \$100 billion.
Suggested Reviewers:	Jochen Hinkel Global Climate Forum hinkel@globalclimateforum.org He is a well known expert on coastal impacts of climate change and also maintains the global DIVA model of coastal effects.

	<p>George Parsons University of Delaware gparsons@udel.edu An economist with expertise in valuation of coastal impacts and beach nourishment as a coastal adaptation technique.</p>
	<p>Dasgupta Susmita Senior Environmental Economist, World Bank sdasgupta@worldbank.org An international expert on the impacts and costs of sea-level rise</p>
	<p>Robin Leichenko Rutgers University robin.leichenko@rutgers.edu Focuses on effects of climate change with a specialization in coastal zones. Serves on editorial board of the Journal of Extreme Events so understands hurricane damage and modeling.</p>

Joint Effects of Storm Surge and Sea-level Rise on US Coasts: New economic estimates of impacts, adaptation, and benefits of mitigation policy

James E. Neumann¹, Kerry Emanuel², Sai Ravela², Lindsay Ludwig¹, Paul Kirshen³, Kirk Bosma⁴, and Jeremy Martinich⁵

Affiliations:

Industrial Economics, 2067 Massachusetts Avenue, Cambridge, MA 02140

WindRiskTech, L.L.C., and Massachusetts Institute of Technology, Cambridge, MA

University of New Hampshire, Durham, NH

Woods Hole Group, Falmouth, MA

USEPA, Climate Change Division, Washington, DC

Abstract:

Recent literature, the US Global Change Research Program's National Climate Assessment, and recent events, such as Hurricane Sandy, highlight the need to take better account of both storm surge and sea-level rise (SLR) in assessing coastal risks of climate change. This study combines three models – a tropical cyclone simulation model; a storm surge model; and a model for economic impact and adaptation – to estimate the joint effects of storm surge and SLR for the US coast through 2100. The model is tested using multiple SLR scenarios, including those incorporating estimates of dynamic ice-sheet melting, two global greenhouse gas (GHG) mitigation policy scenarios, and multiple general circulation model climate sensitivities. The results illustrate that a large area of coastal land and property is at risk of damage from storm surge today; that land area and economic value at risk expands over time as seas rise and as storms become more intense; that adaptation is a cost-effective response to this risk, but residual impacts remain after adaptation measures are in place; that incorporating site-specific episodic storm surge increases national damage estimates by a factor of two relative to SLR-only estimates, with greater impact on the East and Gulf coasts; and that mitigation of GHGs contributes to significant lessening of damages. For a mid-range climate-sensitivity scenario that incorporates dynamic ice sheet melting, the approach yields national estimates of the impacts of storm surge and SLR of \$990 billion through 2100 (net of adaptation, cumulative undiscounted 2005\$); GHG mitigation policy reduces the impacts of the mid-range climate-sensitivity estimates by \$84 to \$100 billion.

1. Introduction

Recent events have caused coastal residents, planners, and government officials to worry about the damaging effects of storm surge, a phenomenon where storm activity can temporarily raise sea levels well beyond the normal tide range. In addition, surging waves associated with these storms have force which can cause great damages to property and infrastructure and put the safety of coastal residents at risk. The study of these effects took on new urgency as a result of the more than \$50 billion in damage caused by Hurricane Sandy, much of it attributed to the effects of storm surge (NCDC 2013, Abel et al. 2012). In recent years, researchers also have begun to link elevated storm surges with sea-level rise (SLR) associated with climate change. For example, an assessment of climate-related risks to the Northeast established that storm surge flood heights which have historically been reached with a 1 in 100 year probability could increase in frequency to rates as high as once every four years, simply because SLR will provide a higher “launch point” for all future storm surges (Frumhoff et al. 2007). That report acknowledged that a quantitative assessment of the likely damages from these events was not yet possible, but recent research has made progress in identifying flood heights from the combined effect of storm surge and SLR over broad areas (Bromirski et al. 2012); in linking models of storm activity to storm surge models for selected areas (Lin et al. 2012); and in conceptualizing the long term damages of storm surge (Kirshen et al. 2012). These effects have been assessed in the coastal chapter of the ongoing US Global Change Research Program’s National Climate Assessment (see Draft for Public Comment as of January 2013), which acknowledges that no national assessment of the joint effect of SLR and storm surge has yet been undertaken.

In this paper, we for the first time extend previous efforts to comprehensively link a tropical cyclone simulation model with a model of storm surge, and assess economic damages, cost-effective adaptation options, and the effects of global greenhouse gas (GHG) mitigation policy using the US Environmental Protection Agency’s (USEPA’s) National Coastal Property Model (NCPM) across the coastline of the contiguous US.

The model is tested using multiple SLR scenarios, including those incorporating estimates of dynamic ice-sheet melting, two GHG mitigation policy scenarios, and multiple general circulation model climate sensitivities. Detailed descriptions of the GHG emissions scenarios,

1
2
3
4 along with projections of global climate change and SLR, are provided in Paltsev et al. (2013). In
5 short, three emission scenarios are used: a reference or ‘business as usual’ (REF), and two
6 scenarios representing futures with policies that limit global GHG emissions such that radiative
7 forcing levels in 2100 are stabilized at 4.5 W/m² (Policy 4.5) or 3.7 W/m² (Policy 3.7). The
8 scenarios used here reflect results for global SLR through 2100, but also incorporate adjustments
9 to account for the omitted effect of dynamic ice-sheet melting, a potentially important factor for
10 SLR projections (Meier et al. 2007). Dynamic ice-sheet melting scenarios incorporate estimates
11 from the empirical model of Vermeer and Rahmstorf (2009), and use as inputs the decadal
12 trajectory of global average air temperature results from the IGSM climate model (Paltsev et al.
13 2013) (see online supplemental material for additional detail on the climate modeling). .
14
15
16
17
18
19
20
21
22
23
24

25 **2. Methods and Data**

26
27
28 The NCPM comprehensively examines the contiguous US coast at a detailed 150m x 150m grid
29 level; incorporates site-specific elevation, land subsidence, and property value data; estimates
30 cost-effective responses to the threat of inundation; and provides economic impact results for
31 three categories of response: shoreline armoring, beach nourishment, and property abandonment
32 (Neumann et al. 2010a and 2010b) (see online supplemental material for additional
33 methodological details for the NCPM). The model was originally developed to address the
34 threat of SLR, and was modified for this work to incorporate the effects of storm surge on
35 estimates of vulnerability, impact, adaptation response, and economic damages.
36
37
38
39
40
41
42
43

44 **Incorporating Storm Surge into the NCPM**

45
46
47 Incorporating storm surge requires modifying the NCPM in three ways: 1) Estimating a
48 cumulative distribution function for location-specific storm surge; 2) Estimating a cumulative
49 distribution function for economic damages; and 3) Adding a new response option (property
50 elevation) that represents a cost-effective alternative in areas subject to episodic flooding but
51 which are not permanently inundated. The first modification relies on the work of Emanuel et al.
52 (2008) for the East and Gulf Coast sites where tropical cyclone activity is the primary threat.
53 Emanuel et al. have developed a method for estimating the generation, track, intensity, and
54 landfall location(s) of simulated tropical cyclone events in climate states represented by large-
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 scale climatology or by global climate models. The wind field output from this simulated storm-
5 generation activity is used as an input in the National Oceanic and Atmospheric Administration's
6 Sea, Lake, and Overland Surge from Hurricanes (SLOSH) model to estimate storm surge depths
7 resulting from these hypothetical storms (Jelesnianski et al, 1992). SLOSH takes into account
8 the storm's pressure, size, forward speed, forecast track, wind speeds, and topographical data.
9 The result is a cumulative density function for storm surge over a century-long time scale at any
10 given location.
11

12
13
14
15
16
17 Tropical cyclones only rarely strike the West Coast, but periodic storms do cause coastal
18 flooding. For the West Coast sites, therefore, we relied on an analysis of historic tide gauge
19 water levels following Tebaldi, Strauss, and Zervas (2012). Hourly and monthly data from the
20 gauges are used to elicit historic patterns of extreme high water events. A peak-over-threshold
21 analysis is performed to select extreme high water events using a threshold corresponding to the
22 99th percentile of each gauge's distribution of observed water levels. The probability of these
23 extreme events is then modeled using a generalized Pareto distribution, the parameters of which
24 can be used to estimate a cumulative density function for storm tide at each gauge. For this
25 analysis, we use SLOSH and the West Coast tide gauge data to model storm surge in the base
26 case (without SLR) over the 21st century. This storm surge cumulative density function is then
27 modified to reflect the effects of SLR on surge height by simply increasing the surge height by
28 the height of relative SLR at any given point in time.
29

30
31
32
33
34
35
36
37
38
39 The second modification to the NCPM follows the approach applied in Kirshen et al. (2012) to
40 estimate a cumulative distribution function for economic damages. First, the storm surge
41 cumulative density function from SLOSH is used to develop an exceedance curve of surge
42 heights. If damages are assigned to each point along the storm surge exceedance curve, then it
43 would become a damage frequency curve. The area under this curve is the annual expected
44 value of storm surge damage. The model calculates damages at eight points along the storm
45 surge exceedance curve (points roughly corresponding to the 2-year, 5-year, 10-year, 20-year,
46 50-year, 100-year, 250-year, and 500-year surge level) using cell-specific data on elevation,
47 property value, and structure and contents depth-damage functions from the U.S. Army Corps
48 (USACE 2000; USACE 2003). These damage values and their corresponding exceedance
49 probabilities are used to estimate the annual expected value of storm surge damage. The storm
50 surge analysis is run on a decadal basis in the NCPM to keep processing time within reasonable
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 limits; therefore, the annual damage estimate is multiplied by ten to estimate damages over the
5 10-year time period. This value is then added to damages calculated in previous time periods to
6 estimate cumulative storm surge damage.
7
8

9
10 The final modification made to the NCPM is the addition of an elevation response option. This
11 response is only allowed as an adaptation to storm surge. We assume a fixed cost to elevate a
12 structure as well as a cost based on amount of fill needed, consistent with the elevation option
13 used in Neumann et al. (2010a).
14
15

16
17 With these modifications, the NCPM is able to calculate storm surge damage for cells at risk.

18
19 Storm surge damages are calculated for all cells with an elevation less than the height of the
20 future (21st century) 500-year storm surge. Note that the effect of adaptation means that some
21 cells with elevation below the 500-year storm surge height are actually not at risk from storm
22 surge, precisely because cells seaward of some vulnerable cells may be modeled to implement a
23 protection response (i.e., armor or nourish), providing protection to the landward cells. The
24 model relies on a “contour analysis” to identify cells that are unprotected and at risk. The
25 contour analysis defines contours based on proximity to open water. Cells adjacent to open
26 water are defined as the first contour. For each cell in this contour, the model first determines
27 whether the cell is at risk. Cells that have armored or nourished are considered “not at risk” from
28 a storm event less than the 100-year surge level (cells protect to withstand a 100-year surge
29 level). Abandoned and elevated cells remain “at risk” from all storm events. Storm surge
30 damage for at-risk cells includes damages from all storms, while storm surge damage for not-at-
31 risk cells includes only damage from storms that “overtop” the protection, or storms greater than
32 the 100-year surge level.
33
34

35
36 The model determines an adaptation response for at-risk cells based on the estimated decadal
37 storm surge damage. In each time period, a cell’s decadal storm surge damage is compared to its
38 property value to determine its response to the threat of storm surge. If the decadal storm surge
39 damage is greater than total property value (structure and land), the cell will be abandoned. If
40 the decadal storm surge damage is less than property value and also less than protection costs,
41 the cell will temporarily incur storm surge damage. If the decadal storm surge damage is less
42 than property value, but greater than protection cost, the property will be protected. The model
43 contains two protection options for storm surge – elevation and armoring (although beach
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 nourishment can also provide some protection from surge). The model chooses the protection
5 option that costs less overall.
6

7
8 After completing the above process for the first contour, the model moves onto cells within the
9 next contour and repeats the process. Cells adjacent to at risk (i.e., unprotected) cells from the
10 first (or previous) contour are considered at risk. Adjacent cells include those in both the
11 cardinal and inter-cardinal directions. The contour analysis continues until all cells have been
12 analyzed.
13
14
15
16

17 **National Extrapolation**

18
19
20 We ran the NCPM to determine the cost of adaptation to and damage from storm surge for 17
21 multi-county study areas on the East coast (Barnstable, MA; New York, NY; Ocean County, NJ;
22 Virginia Beach, VA; Wilmington, NC; Charleston, SC; Jacksonville, FL; and Miami, FL) Gulf
23 coast (Tampa, FL; Pensacola, FL; Mobile, AL; New Orleans, LA; and Galveston, TX), and West
24 coast (Southern California; Northern California; Oregon State; and Washington State). The
25 results of this analysis were extrapolated to the remaining un-modeled coastal counties to
26 calculate a national estimate of tropical cyclone-related costs.
27
28
29

30
31 To accomplish this extrapolation we tried various discrete and continuous methods of assigning
32 to the unmodeled counties a ratio of SLR and storm surge costs to SLR-only costs (note that the
33 SLR-only costs result from running the model without storm surge effects, using the older
34 version of the NCPM described in Neumann et al. 2010). Both types of methods considered
35 factors such as location, area, property value at risk, population, and shoreline length in
36 explaining the results. The continuous approaches attempted did not have much explanatory
37 power, therefore we chose to use a discrete approach of assigning ratios of SLR and storm surge
38 costs to SLR only costs from the modeled counties to the unmodeled counties. The discrete
39 approach used first assigns the unmodeled counties to the closest multi-county study area within
40 the same coastal region (East, Gulf, or West Coast) and then assigns the unmodeled counties to a
41 modeled county within that study area based on the closest match for property value at risk. The
42 ratio of SLR and storm surge costs to SLR only costs, calculated under the REF 3.0°C scenario
43 for the assigned modeled county, is applied to the SLR only results to calculate SLR and storm
44 surge costs for the unmodeled county. The national estimate of tropical cyclone-related costs
45 under the various SLR scenarios is calculated as the sum of storm surge and SLR costs for the
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 counties modeled directly in the NCPM, and storm surge and SLR costs for the counties
5
6 calculated using the extrapolation process described above.
7
8
9

10 **3. Results**

11
12
13
14 Results of our analysis of adaptation to SLR across the seven analyzed scenarios, incorporating
15 dynamic ice sheet melting, are presented in Figure 1. Total undiscounted costs of adaptation to
16 SLR (excluding storm surge) through 2100 range from \$470 billion to \$610 billion under the
17 REF scenarios, depending upon the climate sensitivity value applied, and \$400 billion to \$510
18 billion under the GHG mitigation policy scenarios. Therefore, the benefits of the mitigation
19 scenarios range from \$65 billion to \$98 billion.¹ The largest share of the costs associated with
20 adaptation to SLR are associated with shoreline armoring, followed by nourishment, and the
21 value of abandoned property.
22
23
24
25
26
27

28 As described above, we ran the NCPM to determine the cost of adaptation to and damage from
29 storm surge for 17 multi-county study areas on the East, Gulf, and West Coasts. Figure 2
30 provides the results of this analysis by scenario across the 17 modeled storm surge sites. This
31 figure shows that in the modeled storm surge sites only, the costs associated with storm surge
32 through 2100 range from \$540 billion to \$610 billion under the REF scenarios, depending upon
33 the climate sensitivity value applied, and \$510 billion to \$560 billion under the policy scenarios.
34 Benefits of the GHG mitigation policy scenarios range from \$28 billion to \$49 billion. The
35 largest share of the costs associated with adaptation to SLR and storm surge are associated with
36 shoreline armoring, followed by nourishment, the value of abandoned property, and finally
37 elevating. In addition, a large portion of the costs are associated with residual storm surge
38 damages which occur during storms with surge greater than the height of protection (the NCPM
39 assumes that armoring, nourishment, and elevation is done to protect sufficiently from a 100-
40 year storm) and when the cost-efficient adaptation option is to do nothing and accrue damages.
41 Figure 3 provides a site-specific comparison of the SLR only to the SLR and storm surge results
42 under the REF 3.0°C scenario with dynamic ice sheet melting. Figure 3 also depicts the storm
43
44
45
46
47
48
49
50
51
52
53
54
55
56

57
58 ¹ Discounted at 3% (2005\$ with a 2015 base year), these estimates range from \$200 billion to \$230 billion for the
59 REF scenarios and \$190 billion to \$220 billion for the policy scenarios, thereby yielding mitigation benefits ranging
60 from \$9.4 billion to \$14 billion (see Online Resource 1 for additional detail on discounted results).
61
62
63
64
65

1
2
3
4 surge study areas upon which the national extrapolation was based. Total costs of adaptation to
5 SLR through 2100 range from \$1.3 billion in Washington State to \$51 billion in Miami,
6 increasing to \$1.6 billion in Washington State to \$130 billion in Miami when also considering
7 adaptation to storm surge. In the 17 study areas analyzed, the change in the total cost of
8 adaptation when also considering storm surge, compared to SLR only, ranged from zero percent
9 in Oregon State to an increase of 420 percent in Tampa.

10
11
12
13
14
15 The results of the national extrapolation are provided in Figure 4 for each of the seven scenarios
16 analyzed. Total undiscounted costs of adaptation to SLR and storm surge through 2100 range
17 from \$930 billion to \$1.1 trillion under the REF scenarios, depending upon the climate
18 sensitivity value applied, and \$840 billion to \$980 billion under the different policy
19 scenario/climate sensitivity combinations. This represents an increase of 84 to 110 percent over
20 the costs of adaptation to SLR only. Benefits of the GHG mitigation policy scenarios range from
21 \$84 billion to \$140 billion when considering costs associated with adaptation to both SLR and
22 storm surge.² Figure 5 depicts the cumulative benefits of GHG mitigation over time under the
23 three degree climate sensitivity alternative. This figure shows that benefits do not begin to
24 substantially accrue until mid-century. Beginning around 2050, benefits accrue rapidly and at a
25 more substantial pace when also considering impacts associated with storm surge.

36 37 38 **4. Discussion and Conclusions**

39
40
41 The results presented in this paper indicate the importance of considering the combined effect of
42 SLR and storm surge when analyzing the risk of climate change to coastal property.

43
44
45 Incorporating episodic storm surge increases the undiscounted national scale costs associated
46 with adaptation by approximately a factor of two, compared to an SLR only estimate. Further,
47 GHG mitigation results in cumulative undiscounted avoided costs of up to \$98 billion through
48 2100 when considering SLR alone and up to \$140 billion when considering the joint effect of
49 SLR and storm surge. Because GHG mitigation has a delayed effect on SLR, the majority of the

50
51
52
53
54
55
56
57
58 ² Discounted at 3% (2005\$ with a 2015 base year), these estimates range from \$680 billion to \$730 billion for the
59 REF scenarios and \$660 billion to \$700 billion for the policy scenarios, thereby yielding mitigation benefits ranging
60 from \$18 billion to \$29 billion (see online supplemental material for additional detail on discounted results).
61
62
63
64
65

1
2
3
4 benefits associated with mitigation are not felt until mid-century. Benefits increase rapidly
5
6 beginning around 2050.

7
8 While mitigation does have a substantial effect on reducing future damages of climate change on
9
10 coastal property, adaptation plays a crucial role in responding to the threat. In particular, due to
11
12 the high value of coastal property, driven by the high amenity value of coastal property, the
13
14 optimal adaptation response more often involves protection rather than abandonment of property.
15
16 When considering SLR only, costs associated with armoring and beach nourishment represent
17
18 approximately 59 and 29 percent of the total costs associated with adaptation, respectively.
19
20 Abandonment of property makes up the remaining 12 percent of these costs. When also
21
22 considering storm surge in the 17 modeled sites, costs associated with armoring, nourishment,
23
24 and elevation represent a smaller share of the costs associated with adaptation, between 66 and
25
26 74 percent. Abandonment becomes an optimal adaptation option in more cases due to the
27
28 significant damages caused by storm surge. This is illustrated in Figure 6 for the Tampa study
29
30 site – the left panels illustrate the cost-effective adaptation response to SLR risks, with red areas
31
32 indicating abandonment, black areas lines of armoring defense, yellow areas beach nourishment,
33
34 and brown areas structure elevation. The incremental effect of dynamic ice sheet melting is
35
36 shown in the bottom panels. A significant difference in abandoned area is evident when
37
38 comparing the left and right panels. In particular, the red areas in the low elevation east bay
39
40 show great sensitivity to storm surge. A similar map for New York City would show less
41
42 abandonment and more protection and elevation in response to risks of episodic flooding, owing
43
44 to higher property values in New York City’s vulnerable areas.

45
46 It is also important to note that there are significant residual impacts due to storm surge after
47
48 adaptation measures are in place. These damages make up between 25 and 31 percent of the
49
50 total costs associated with SLR and storm surge in the modeled sites. The residual impacts
51
52 include two components. The first component arises from areas where episodic flooding occurs,
53
54 but where the extent and damage from flooding are not so great as to trigger an abandonment
55
56 response. The second component arises in areas where protection is warranted, but there is
57
58 infrequent overtopping of the protection for rare but severe storms. The first component
59
60 accounts for the largest portion of residual impacts, on an expected value basis, but the second
61
62 component, associated with severe storms, may underestimate effects because of our current
63
64
65

1
2
3
4 inability to estimate indirect effects such as loss of critical infrastructure, business interruption,
5 and debris removal costs that may be a much larger component of damages than the direct
6 structure damage in these more severe storms.
7
8
9

10
11 The effect of storm surge varies dramatically by location. As expected, costs associated with
12 storm surge vary geographically due to topography, value of coastal property, and the nature of
13 storms. In some areas, responding to the gradual effects of SLR provides nearly sufficient
14 protection from storm surge as well. But in many other areas, storm surge is significant
15 amplifying factor. Low lying areas with significant coastal development that have greater storm
16 activity have the greatest costs associated with storm surge. Under the REF scenario with 3°C
17 climate sensitivity, the four sites with the greatest increase in costs when also considering storm
18 surge are Tampa (420 percent increase), New York City (220 percent increase), New Orleans
19 (210 percent increase), and Galveston (190 percent increase). In recent years major storms have
20 hit the New York, New Orleans, and Galveston areas showing firsthand how susceptible these
21 areas are to storm surge.
22
23
24
25
26
27
28
29
30

31 The approach presented in this paper does have some limitations. For example, we are currently
32 unable to capture damages from wind or rain associated with coastal storms. The spatial extent
33 as well as the magnitude of the damages are likely to increase when also considering the effect of
34 wind and rain (see Mendelsohn et al. 2012 for more details on the potential for wind damage
35 from tropical cyclones). In addition, our approach is designed to analyze impacts to coastal
36 property, but does not explicitly account for impacts to infrastructure and ecosystem services.
37 The model also does not include post-disaster damages such as those from power outages and
38 lost productivity (Abel et al. 2012). We recognize that the effect of a major storm on the
39 economy of a region may be felt long after the actual storm. In addition, for the East and Gulf
40 coasts, our analysis focuses on storms of tropical origin. Extra-tropical storms also cause
41 significant damage but are not addressed here (see Narita et al. 2010 for more information).
42
43
44
45
46
47
48
49
50
51
52

53 Three key directions emerge for further research. First, while the sea-level rise impacts and
54 adaptation literature has firmly established that adaptation such as that modeled here is a very
55 cost-effective response to coastal inundation and flood risks, recent storm events have shown that
56 in many cases cost-effective adaptation measures have yet to be adopted. This suggests that
57
58
59
60
61
62
63
64
65

1
2
3
4 further attention should be paid to establishing the level of economic assets that are vulnerable to
5 the coastal SLR and storm surge risks modeled here, absent adaptation, and evaluating how
6 incremental “tiers” of adaptation action can serve to mitigate those risks, with costs and benefits
7 tracked along the way. Second, for computational reasons we focus our national estimates on the
8 application of the SLOSH storm surge model – but in many locations a more detailed
9 hydrodynamic model (e.g., ADCIRC) that includes all key physical processes (e.g., wind, waves,
10 freshwater discharge) is needed to better characterize surge risks. Further site-specific
11 comparisons are needed along the lines of Lin et al. (2012) that include evaluation of the impact
12 on economic assessments such as ours when using these more refined storm surge tools. In
13 addition, in areas identified at high risk from storm surge likely justify a modest analytic
14 investment in ADCIRC modeling, in part to provide better assurance that planned adaptations to
15 storm surge can be both effective and cost-effective, before the much larger infrastructure
16 investments are made. Third, recent research suggests that SLR is likely to manifest with
17 regional “hotspots” of unusually high SLR. One of these hotspots is the US North Atlantic coast,
18 an area with substantial economic value near the coast (Sallenger et al. 2012). Future analyses
19 should take into account these emerging findings and test the sensitivity of impact and adaptation
20 cost results to higher levels of SLR that reflect the non-uniform influence of currents and other
21 local factors.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 Acknowledgement: The authors wish to acknowledge the financial support of the USEPA’s Climate Change
43 Division (Contracts EP-D-09-054 and EP-BPA-12-H-0024). The views expressed in this document do not
44 necessarily represent those of the USEPA. In addition, we are grateful to Benjamin Strauss of Climate Central for
45 providing historic storm data for selected West Coast sites.
46
47
48
49
50

51 **References**

52
53
54 Abel, J R, Bram J, Deitz R, and Orr J. (2012) What Are the Costs of Superstorm Sandy? Federal Reserve Bank of
55 New York. Available online at [http://libertystreeteconomics.newyorkfed.org/2012/12/what-are-the-costs-of-](http://libertystreeteconomics.newyorkfed.org/2012/12/what-are-the-costs-of-superstorm-sandy.html)
56 [superstorm-sandy.html](http://libertystreeteconomics.newyorkfed.org/2012/12/what-are-the-costs-of-superstorm-sandy.html)
57
58 Bromirski PD, Cayan DR, Graham N, Flick RE, Tyree M (Scripps Institution of Oceanography) (2012) Coastal
59 Flooding Potential Projections: 2000–2100. California Energy Commission, CEC-500-2012-XXX
60
61
62
63
64
65

1
2
3
4
5 Emanuel K, Sundararajan R, Williams J (2008) Hurricanes and Global Warming: Results from Downscaling IPCC
6 AR4 Simulations. *Bulletin of the American Meteorological Society* 89: 347–67
7
8
9

10 Frumhoff PC, McCarthy JJ, Melillo JM, Moser SC, Wuebbles DJ (2007) *Confronting Climate Change in the U.S.*
11 *Northeast: Science, Impacts, and Solutions*. Synthesis report of the Northeast Climate Impacts Assessment (NECIA).
12 Cambridge, MA: Union of Concerned Scientists (UCS)
13
14

15
16 Jelesnianski, CP, Chen J, and Shaffer WA (1992). SLOSH: Sea, lake, and overland surges from hurricanes. NOAA
17 Technical Report NWS 48, National Oceanic and Atmospheric Administration, U. S. Department of Commerce,
18 Washington, DC.
19
20
21

22 Kirshen P, Merrill S, Slovinsky P, and Richardson N (2012) Simplified Method for Scenario-Based Risk
23 Assessment Adaptation Planning in the Coastal Zone. *Climatic Change*, 113: 3-4. doi: 10.1007/s10584-011-0379-z
24
25

26
27 Lin N, Emanuel K, Oppenheimer M, Vanmarcke E (2012) Physically based assessment of hurricane surge threat
28 under climate change. *Nature Climate Change* 2: 462–467. doi:10.1038/nclimate1389
29
30

31 Meier MF, Dyurgerov MB, Rick UK, O’Neel S, Pfeffer WT, Anderson RS, Anderson SP, Glazovsky AF (2007)
32 *Glaciers Dominate Eustatic Sea-Level Rise in the 21st Century*. *Science* 317: 1064–1067
33
34
35

36 Mendelsohn R, Emanuel K, Chonabayashi S, Bakkensen L (2012) The impact of climate change on global tropical
37 cyclone damage. *Nature Climate Change* 2:205–209. doi:10.1038/nclimate1357
38
39

40 Narita D, Tol R, Anthoff D (2010) Economic costs of extratropical storms under climate change: An application of
41 FUND. *Journal of Environmental Planning and Management*, 53 (3). pp. 371-384
42
43
44

45 NCDC (National Climatic Data Center) (2013) *Billion-Dollar U.S. Weather/Climate Disasters 1980-2013*
46 National Climatic Data Center, Asheville, NC. Downloaded from <http://www.ncdc.noaa.gov/billions/>
47
48
49

50 Neumann JE, Hudgens DE, Herter J, Martinich J (2010a) Assessing sea-level rise impacts: a GIS-based framework
51 and application to coastal New Jersey. *Coastal Management* 38(4): 433-455
52
53

54 Neumann JE, Hudgens DE, Herter J, Martinich J (2010b) *The Economics of Adaptation along Developed*
55 *Coastlines*. *Wiley Interdisciplinary Reviews: Climate Change* 2(1): 89-98
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Paltsev S, Monier E, Scott J, Sokolov A, Reilly J, Gao X, Schlosser A (2013) Integrated economic and climate projections for impact assessment. *Clim Change*. doi: 10.1007/s10584-013-0892-3

Sallenger, A.H., K.S. Doran and P.A. Howd (2012) Hotspot of accelerated sea-level rise on the Atlantic coast of North America. *Nature Climate Change* 2: 884-888


Tebaldi C, Strauss B, Zervas C (2012) Modeling sea-level rise impacts on storm surges along US coasts *Environ. Res. Lett.* 7 014032

USACE (2000) Economic Guidance Memorandum (EGM) 01-03, Generic Depth-Damage Relationships. 4 Dec 2000


USACE (2003) Economic Guidance Memorandum (EGM) 04-01, Generic Depth-Damage Relationships for Residential Structures with Basements. 10 October 2003

Vermeer M, Rahmstorf S (2009) Global sea level linked to global temperature. *Proceedings of the National Academy of Sciences* 106(51): 21527-21532

1
2
3
4 Fig. 1 Cumulative total costs of adaptation in the contiguous US by 2100 for sea-level rise with
5 dynamic ice sheet melting, undiscounted (\$ million). CS is climate sensitivity of the underlying
6 general climate model
7
8
9



10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34 Fig. 2 Cumulative total costs of adaptation by 2100 for sea-level rise with dynamic ice sheet
35 melting and storm surge for the 17 modeled sites only, undiscounted (\$ million)
36
37



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Fig. 3 Total costs of adaptation by 2100 under reference 3° scenario with dynamic ice sheet melting, undiscounted (\$ million)




Fig. 4 Comparison of cumulative total costs of adaptation by 2100 in the contiguous US for sea-level rise with dynamic ice sheet melting with and without storm surge, undiscounted (\$ million)






Fig. 5 Cumulative benefits in the contiguous US for sea-level rise with dynamic ice sheet melting with (dashed line) and without (solid line) storm surge, undiscounted (\$ millions)



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig. 6 Effect of incorporating storm surge in economic impact estimates for Tampa, Florida
(NOTE: better quality graphic available for publication purposes)



1
2
3
4 **Online Resource 1**
5

6
7 **Joint Effects of Storm Surge and Sea-level Rise on US**
8 **Coasts: New economic estimates of impacts, adaptation, and**
9 **benefits of mitigation policy**
10
11
12
13
14

15
16
17 James E. Neumann¹, Kerry Emanuel², Sai Ravela², Lindsay Ludwig¹, Paul Kirshen³, Kirk
18 Bosma⁴, and Jeremy Martinich⁵
19
20

21
22
23 *Affiliations:*
24

25 *Industrial Economics, 2067 Massachusetts Avenue, Cambridge, MA 02140*

26 *WindRiskTech, L.L.C., and Massachusetts Institute of Technology, Cambridge, MA*

27 *University of New Hampshire, Durham, NH*
28

29 *Woods Hole Group, Falmouth, MA*

30 *USEPA, Climate Change Division, Washington, DC*
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig. 1 Sea-level rise with dynamic ice sheet melting




Fig. 2 Cumulative total costs of adaptation by 2100 in the contiguous US for sea-level rise with dynamic ice sheet melting, discounted at 3% (\$ million)






Fig. 3 Cumulative total costs of adaptation by 2100 for sea-level rise with dynamic ice sheet melting and storm surge for the 17 modeled sites only, discounted at 3% (\$ million)



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Fig. 4 Total costs of adaptation by 2100 under reference 3° scenario with dynamic ice sheet melting, discounted at 3% (\$ million)



Basemap Sources:
USGS, Esri, TANA, AND

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig. 5 Comparison of cumulative total costs of adaptation by 2100 in the contiguous US for sea-level rise with dynamic ice sheet melting with and without storm surge, discounted at 3% (\$ million)





Fig. 6 Cumulative benefits in the contiguous US for sea-level rise with dynamic ice sheet melting with (dashed line) and without (solid line) storm surge, discounted at 3% (\$ millions)



Additional Information on Climate Projections

Three emissions scenarios are used: a reference (REF) or business as usual scenario, corresponding to a radiative forcing of 9.8 W/m^2 by 2100, and two future scenarios that limit global GHG emissions such that the global radiative forcing levels in 2100 are stabilized at 3.7 W/m^2 (Policy 3.7) and 4.5 W/m^2 (Policy 4.5). In addition, three alternative climate sensitivity values are applied in the analyses of this paper: 2°C , 3°C and 6°C .

Using these scenarios, future climate projections for projecting SLR effort were developed using the National Center for Atmospheric Research's Community Atmospheric Model (Version 3), a three-dimensional atmosphere model, linked with the Integrated Global Systems Model (IGSM-CAM) (Monier et al. 2013). As a full general circulation model (GCM), IGSM-CAM generates projections that incorporate not only changes in mean climate, but also incorporates features explicitly designed to allow for year-to-year variability. Decadal SLR projections (absent dynamic ice-sheet melting) were a direct output of the IGSM-CAM model. As described in the text, estimates that include dynamic ice sheet melting were derived from the empirical model of Vermeer and Rahmstorf (2009) using average ambient air temperature trajectories from the IGSM-CAM model. Storm surge estimates derive from hurricane modeling that simulates storm generation activity over the 20st century using backcasting results from the GFDL Climate Model for the historic period of 1956 to 2005.

References

Monier E, Gao X, Scott J, Sokolov A, Schlosser A (2013) A framework for modeling uncertainty in regional climate change. *Climatic Change*


Vermeer M, Rahmstorf S (2009) Global sea level linked to global temperature. *Proceedings of the National Academy of Sciences* 106(51): 21527-21532

Additional Methodological Details for EPA’s National Coastal Property Model (NCPM)

The overall framework of the model is presented in Figure 1. The basic structure involves arraying relevant input data, listed on the left side of Figure 1, and constructing a spatial geodatabase on a 150-m grid cell frame, which can then be analyzed to estimate response to SLR, the cost of the adaptive response, and the ‘residual damages’ that result in areas where adaptive measures are not cost-effective, and therefore flooding damages are evident. The grid frame encompasses virtually all areas potentially vulnerable to the effects of SLR and storm surge, including approximately 300 coastal counties in the continental US. Most of these counties have direct coastal or bay frontage, but some are affected only through proximity to tidally influenced rivers and tributaries, a common geographic feature in the Southern Atlantic and Gulf regions. Analysis and aggregation modules of the model, depicted in the center right and bottom right corner of Figure 1, access the site-specific data within the geodatabase (e.g., elevation) along with a series of user-defined input parameters (e.g., SLR scenario and the cost of armoring or beach nourishment), and estimate the timing and costs of adapting to SLR over time. Armoring involves a hard structure of some sort—prior work and review suggests concrete structures on ocean-fronting areas, and less expensive bulkheads on bay-fronting areas—and beach nourishment involves placing sand on beach areas. In all cases, long-term maintenance of structures or period re-nourishment is also required, with differential costs for each.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1:



Some elements of the framework, particularly the optimal response model, rely extensively on earlier research by Yohe et al. (1996 and 2011) assessing economic impacts for 30 coastal US sites. The optimal response approach is based on a simplified benefit–cost analysis of protective structures and beach nourishment relative to a retreat response. The planning problem expressed as an optimization problem involves choosing a time between today and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2100 to initiate a protection plan as adaptation to SLR which maximizes the following expression:

$$PV\{B[t_0,T]\} - PV\{C[t_0,T]\} \quad (1)$$

where t_0 is the time when protection as adaptation is initiated, T is the time when protection that had been initiated subsequently might be abandoned, and the present value of B and C are expressions of the benefits and costs, respectively, of protection over time as a function of the choice of t_0 . Prior theoretical work established that if hard structure protection prior to 2100 is optimal, then it remains optimal to continue to maintain that hard structure in perpetuity, making the otherwise difficult estimation of T irrelevant. As a result, this generic model simplifies to the following decision rules. First, where the cost of measures designed to protect properties from SLR is less than the benefit of avoided property value loss, the time to begin protection that maximizes Eq. (1) is earlier than 2100, and the adaptation cost incurred in response to SLR is estimated as the capital cost of construction plus ongoing maintenance costs. Second, where the estimated protection cost exceeds the benefit of adaptation, the expression is maximized when t_0 is equal to the time horizon of the simulation, in this case 2100. Then retreat (i.e., abandonment) is the estimated response to the threat, and the impact of SLR is lost structure and land value.

Projecting resources at risk involves, at minimum, estimating how real property values could appreciate—our approach links future property value to a projection of US gross domestic product (GDP) developed as part of a broader EPA project (see Neumann et al. 2014).

Implicit in our response analysis are two key features of the approach: (1) we estimate the optimal timing of a response, largely based on the timing of inundation and (2) abandonment decisions are irreversible, and protection and beach nourishment decisions, while theoretically reversible, are also effectively permanent. Estimating optimal timing demands that we rely on an SLR trajectory, rather than simply an endpoint, and also critically affects the economic cost calculations through application of a positive discount rate. Irreversible decisions are not a requirement of the approach, but prior work suggests that, at least within the optimal response paradigm, these decisions are made once and remain robust over time.

References

1 Neumann JE, Price J, Chinowsky P, Wright L, Ludwig L, Streeter R, Jones R, Smith JB,
2 Perkins W, Jantarasami L, and Martinich J. (2014) Climate change risks to US
3 infrastructure: impacts on roads, bridges, coastal development, and urban drainage. Climatic
4 Change, DOI 10.1007/s10584-013-1037-4.
5
6
7
8
9

10 Yohe, G, Neumann, JE, Marshall, P and Ameden, H. (1996) The Economic Cost of
11 Greenhouse Induced Sea Level Rise in the United States. Climatic Change. 32: 387-410.
12
13
14
15

16 Yohe, G, Knee, K, and Kirshen, P (2011) On the economics of coastal adaptation solutions in
17 an uncertain world, Climatic Change, 106:71–92.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

Abel, J R, Bram J, Deitz R, and Orr J. (2012) What Are the Costs of Superstorm Sandy? Federal Reserve Bank of New York. Available online at <http://libertystreeteconomics.newyorkfed.org/2012/12/what-are-the-costs-of-superstorm-sandy.html>

Bromirski PD, Cayan DR, Graham N, Flick RE, Tyree M (Scripps Institution of Oceanography) (2012) Coastal Flooding Potential Projections: 2000–2100. California Energy Commission, CEC-500-2012-XXX

Emanuel K, Sundararajan R, Williams J (2008) Hurricanes and Global Warming: Results from Downscaling IPCC AR4 Simulations. *Bulletin of the American Meteorological Society* 89: 347–67

Frumhoff PC, McCarthy JJ, Melillo JM, Moser SC, Wuebbles DJ (2007) *Confronting Climate Change in the U.S. Northeast: Science, Impacts, and Solutions*. Synthesis report of the Northeast Climate Impacts Assessment (NECIA). Cambridge, MA: Union of Concerned Scientists (UCS)

Jelesnianski, CP, Chen J, and Shaffer WA (1992). SLOSH: Sea, lake, and overland surges from hurricanes. NOAA Technical Report NWS 48, National Oceanic and Atmospheric Administration, U. S. Department of Commerce, Washington, DC.

Kirshen P, Merrill S, Slovinsky P, and Richardson N (2012) Simplified Method for Scenario-Based Risk Assessment Adaptation Planning in the Coastal Zone. *Climatic Change*, 113: 3-4. doi: 10.1007/s10584-011-0379-z

Lin N, Emanuel K, Oppenheimer M, Vanmarcke E (2012) Physically based assessment of hurricane surge threat under climate change. *Nature Climate Change* 2: 462–467. doi:10.1038/nclimate1389

Meier MF, Dyurgerov MB, Rick UK, O’Neel S, Pfeffer WT, Anderson RS, Anderson SP, Glazovsky AF (2007) Glaciers Dominate Eustatic Sea-Level Rise in the 21st Century. *Science* 317: 1064–1067

Mendelsohn R, Emanuel K, Chonabayashi S, Bakkensen L (2012) The impact of climate change on global tropical cyclone damage. *Nature Climate Change* 2:205–209. doi:10.1038/nclimate1357

Narita D, Tol R, Anthoff D (2010) Economic costs of extratropical storms under climate change: An application of FUND. *Journal of Environmental Planning and Management*, 53 (3). pp. 371-384

NCDC (National Climatic Data Center) (2013) Billion-Dollar U.S. Weather/Climate Disasters 1980-2013 National Climatic Data Center, Asheville, NC. Downloaded from <http://www.ncdc.noaa.gov/billions/>

Neumann JE, Hudgens DE, Herter J, Martinich J (2010a) Assessing sea-level rise impacts: a GIS-based framework and application to coastal New Jersey. *Coastal Management* 38(4): 433-455

Neumann JE, Hudgens DE, Herter J, Martinich J (2010b) The Economics of Adaptation along Developed Coastlines. *Wiley Interdisciplinary Reviews: Climate Change* 2(1): 89-98

Paltsev S, Monier E, Scott J, Sokolov A, Reilly J, Gao X, Schlosser A (2013) Integrated economic and climate projections for impact assessment. *Clim Change*. doi: 10.1007/s10584-013-0892-3

Sallenger, A.H., K.S. Doran and P.A. Howd (2012) Hotspot of accelerated sea-level rise on the Atlantic coast of North America. *Nature Climate Change* 2: 884-888

Tebaldi C, Strauss B, Zervas C (2012) Modeling sea-level rise impacts on storm surges along US coasts *Environ. Res. Lett.* 7 014032

USACE (2000) Economic Guidance Memorandum (EGM) 01-03, Generic Depth-Damage Relationships. 4 Dec 2000

USACE (2003) Economic Guidance Memorandum (EGM) 04-01, Generic Depth-Damage Relationships for Residential Structures with Basements. 10 October 2003

Vermeer M, Rahmstorf S (2009) Global sea level linked to global temperature. *Proceedings of the National Academy of Sciences* 106(51): 21527-21532